


State of California Sea Level Rise Guidance

2024 SCIENCE & POLICY UPDATE



About the Report

THIS REPORT WAS PRODUCED by the California Ocean Protection Council (OPC), in partnership with the California Ocean Science Trust (OST) and a scientific Task Force (Task Force). This report updates and replaces the previous 2018 ‘State of California Sea-Level Rise Guidance’ and marks the fourth iteration of statewide guidance since 2010 for state and local decision-makers to incorporate best available science on sea level rise into planning, design, permitting, investments, and other decisions. To support science-based decision-making, this report consists of syntheses of the best available science on sea level rise and other coastal hazards (e.g., flooding and erosion) with pragmatic and practical approaches for using this new scientific information (Chapters 2.0 and 3.0), primarily led and authored by the Task Force, as well as specific policy recommendations for incorporating this information into decision-making (Chapter 4.0), led and authored by OPC. To ensure this report meets the diversity of needs and interests for sea level rise decision-making and planning across California, state and local decision-makers, California Native American tribes, planners, and other practitioners were consulted throughout to provide input and feedback into the process and content of this report. An external scientific panel of peer reviewers provided critical review of the sea level rise science synthesis. To keep pace with future advancements in scientific understanding of sea level rise, OPC will continue to uphold its commitment to update this statewide guidance approximately every five years.

CALIFORNIA OCEAN PROTECTION COUNCIL

OPC is a Cabinet-level state body that works jointly with state and federal agencies, non-governmental organizations (NGOs), tribes and the public to ensure that California maintains healthy, resilient, and productive ocean and coastal ecosystems. They do so by advancing innovative science-based policy and management, making strategic investments and catalyzing action through partnerships and collaboration. OPC commissioned the science update and authored the policy guidance (Chapter 4) for application in planning and project decisions.

CALIFORNIA OCEAN SCIENCE TRUST

OST is an independent non-profit created by the California Legislature to bring cutting edge science to the decisions shaping the future of the California coast and ocean. OST led the delivery of sound, policy-relevant scientific advice to support the development of this report and meet the needs of state and local decision-makers. This included establishing the vision for this effort with the OPC, designing the collaborative science-policy process, and convening the Task Force.



SEA LEVEL RISE SCIENCE TASK FORCE

The Sea Level Rise Task Force is an interdisciplinary and multi-institutional collaborative of scientific experts convened beginning in June 2022 by OST to update California’s sea level rise guidance based on recent scientific advancements. Members of the Task Force contributed their scientific expertise and perspectives throughout the process to the technical scientific chapters of this report (Chapters 2.0 and 3.0) but did not provide policy direction. Authorship is noted at the top of each chapter to reflect contributions accordingly.

Dr. Susheel Adusumilli

University of California, San Diego

Dr. Patrick Barnard

*U.S. Geological Survey**

Dr. Daniel Cayan

University of California, San Diego

Laura Engeman

*California Sea Grant & University of California, San Diego**

Dr. Gary Griggs

University of California, Santa Cruz

Dr. Benjamin Hamlington

*National Aeronautics and Space Administration**

Dr. Kristina Hill

University of California, Berkeley

Dr. Felix Landerer

National Aeronautics and Space Administration

Dr. Phil Thompson

University of Hawaii at Manoa

**Task Force Co-Chairs*

SUGGESTED CITATION

California Sea Level Rise Guidance: 2024 Science and Policy Update. 2024. California Sea Level Rise Science Task Force, California Ocean Protection Council, California Ocean Science Trust.

SUGGESTED IN-TEXT CITATION

California Sea Level Rise Guidance (2024)

ACKNOWLEDGEMENTS

We are grateful to the agency and departments in the State Sea Level Rise Collaborative, California Natural Resources Agency and the local partners who comprised our Sea Level Rise Advisory Panel including David Behar, Dana Brechwald, Dr. Charles Lester, Hilary Papendick, and Dr. Laurie Richmond who provided meaningful input and feedback on this report, as well as the external peer reviewers whose expertise assisted in ensuring this report is based on credible science: Dr. Robert Kopp, Dr. Sophie Crooks Nowicki, Dr. Eric Rignot, and Dr. William Sweet.

Table of Contents

| | |
|---|-----------|
| EXECUTIVE SUMMARY | 5 |
| Why Are We Updating the Guidance | 5 |
| Statewide Averages for Five California Sea Level Scenarios | 7 |
| Key Takeaways | 8 |
| 1. INTRODUCTION | 10 |
| 1.1. Existing State Policy Guidance and Sea Level Rise Planning | 11 |
| 1.2. Purpose & Intended Use | 13 |
| 1.3. How This Report was Developed | 14 |
| 1.4. Updating the Science Foundation for Statewide Policy Guidance | 14 |
| 1.5. Translating Science Into Policy Guidance | 16 |
| 2. CALIFORNIA SEA LEVEL SCENARIOS | 19 |
| 2.1. Updated Scientific Understanding | 20 |
| 2.2. Steps to Build California Sea Level Scenarios | 23 |
| 2.3. Sea Level Scenarios 2020-2150 | 26 |
| 2.4. Sea Level Scenario Storylines | 34 |
| 3. COMBINED IMPACTS OF SEA LEVEL RISE AND OTHER COASTAL HAZARDS | 38 |
| 3.1. Flooding and Sea Level Rise | 39 |
| 3.2. Groundwater and Sea Level Rise | 44 |
| 3.3. Coastal and Shoreline Erosion | 48 |
| 3.4. Preparing for Extreme Coastal Storms | 52 |
| 4. CALIFORNIA SEA LEVEL RISE POLICY GUIDANCE | 53 |
| 4.1. Summary | 53 |
| 4.2. Stepwise Process to Apply Sea Level Scenarios in Planning and Projects | 54 |
| 4.3. General recommendations for Sea level Rise Planning and Adaptation | 67 |
| 5. CONCLUSIONS AND LOOKING FORWARD | 72 |
| 6. REFERENCES | 73 |
| APPENDICES | |
| Appendix 1. Map of California NOAA Tide Gauge Locations (14 total) | 84 |
| Appendix 2. Sea Level Scenarios at NOAA Tide Gauge Locations | 85 |
| Appendix 3. Localized Assessment of Vertical Land Motion (VLM) and Incorporation into Sea Level Scenarios | 93 |
| Appendix 4. Tools and Resources to Support Visualization of Sea Level Rise and Coastal Hazards | 97 |
| Appendix 5. Technical Detail on Formation of Sea Level Scenarios | 100 |

Executive Summary

AUTHORS: Ocean Protection Council and Ocean Science Trust

CLIMATE CHANGE is altering California's coastline. Rising seas, colliding with more frequent and extreme storms, are drowning beaches, eroding bluffs, flooding homes and businesses, and damaging roads and other essential public infrastructure. Close to 70% of California's residents live in coastal counties, and millions more visit every year, driving the state's \$44 billion-dollar coastal economy¹ as people come to the coast for recreation, cultural and spiritual well-being, a connection to nature, and to support their livelihoods.

To ensure that people, the environment, and the economy can continue to thrive, California must take bold and swift action to help prepare communities for the impacts we are starting to see now and that are projected to worsen in the years ahead. The Newsom Administration and the State Legislature have taken this responsibility seriously, demonstrating leadership by passing landmark legislation, providing unprecedented funding for coastal resilience planning and adaptation projects, including large-scale restoration efforts, and creating a State Sea Level Rise Action Plan to align agency priorities and decisions and leverage expertise and resources across California.

Failure to adequately prepare now will have significant cost implications in the future and consequences to public health and safety, wildlife and habitats, private property, water supply, and infrastructure necessary to maintain daily living in California. It will also have impacts on communities burdened by social and environmental injustice who are already disproportionately impacted by climate change, industrialization, and pollution.

To build resilience for coastal communities and ecosystems, thoughtful science-based planning and adaptation actions need to happen now. This updated State of California

Sea Level Rise Guidance provides the best available science and policy recommendations from which to make these decisions. California's enduring connection to the coast demands that we acknowledge the threats on the horizon and innovate to adapt to the changes ahead.

Why Are We Updating the Guidance

The previous State of California Sea-Level Rise Guidance was issued in 2018 (referred to hereafter as the 2018 California Sea-Level Rise Guidance) and was based on a synthesis of best available science at that time. Since then, there have been significant advancements in scientific understanding and ability to project future sea level rise. In February 2022 a national report entitled Global and Regional Sea Level Rise Scenarios for the United States (referred to hereafter as the 2022 Federal Sea Level Rise Technical Report) was released updating Sea Level Scenarios for the United States based on global projections in the latest Intergovernmental Panel on Climate Change report. This national update presented an opportunity to update California's sea level rise guidance with best available science and align the state's approach with national coastal adaptation efforts.

This report, which includes updated Sea Level Scenarios and policy recommendations, serves as the 2024 update to the 2018 California Sea-Level Rise Guidance. It will support state and local action to assess vulnerability to rising seas and climate-driven flooding and the creation of adaptation plans and projects that build resilience into the future.

1. <https://coast.noaa.gov/data/digitalcoast/pdf/california-ocean-economy.pdf>

BOX 1:**California Sea Level Scenarios**

Five Sea Level Scenarios are constructed and presented for California. Adopting the scientific framework and approach used in the 2022 Federal Sea Level Rise Technical Report and creating consistency between state and federal planning, each scenario is defined and labeled according to a target value of global mean sea level rise in 2100 (e.g., the Intermediate Scenario has a GMSL target of 1.0m (3.3ft)). The Sea Level Scenarios are derived from the sets of probabilistic projections developed in the Intergovernmental Panel on Climate Change Sixth Assessment report (IPCC AR6) and reflect the most up to date scientific understanding of the physical drivers of sea level rise. The information about the likelihood of meeting or exceeding a specific Sea Level Scenario is embedded in the scenarios themselves (e.g., the High Scenario is less likely than the Intermediate Scenario, as described in more detail in Chapter 2.0). The Sea Level Scenarios for California span the plausible range of future sea level rise under all emissions and global development futures and enable users to consider sea level rise without first selecting a single emissions future on which to base planning and projects.

LOW: 0.3m (1.0ft) by 2100 - The target of 1 foot of increase in global sea level rise by 2100 is set under the assumption of the current rate of sea level rise continuing on into the future. This assumption is inconsistent with current observations of an acceleration in sea level rise, but could still be considered plausible under the most aggressive emission reduction scenarios. This scenario is on the lower bounding edge of plausibility given current warming and sea level trajectories.

INTERMEDIATE-LOW: 0.5m (1.6ft) by 2100 - This scenario arises under a range of future emissions pathways, associated with a range of future warming levels and socioeconomic development pathways. Given current sea level observations and estimates of future warming, this scenario provides a reasonable estimate of the lower bound for the most likely sea level rise by 2100. Since low confidence processes (e.g., rapid ice sheet melt) are not important to this scenario, the range of possible sea level rise after 2100 does not expand significantly.

INTERMEDIATE: 1.0m (3.3ft) by 2100 - This scenario is driven dominantly by high emissions scenarios, and thus higher warming levels.

Projections including contributions from low confidence processes provide about 25% of the pathways for reaching the scenario target by 2100. This scenario provides a reasonable upper bound for the most likely range of sea level rise by 2100.

INTERMEDIATE-HIGH: 1.5m (4.9ft) by 2100 - This scenario is heavily reflective of a world where rapid ice sheet loss processes are contributing to sea level rise, associated with intermediate to high future emissions, and high warming. The amount of sea level rise by 2100 corresponds to other scientific estimates of plausible high-end projections.

HIGH: 2.0m (6.6ft) by 2100 - This scenario only arises with high future emissions and high warming with large potential contributions from rapid ice sheet loss processes. Deep uncertainties and ambiguity embedded in this scenario frame a worst case beyond 2100 as we currently understand it, and a statement about the likelihood of reaching this scenario is not possible. It should be used with caution and consideration of the underlying assumptions.

Statewide Averages for Five California Sea Level Scenarios

Median values for California Sea Level Scenarios, in feet, relative to a 2000 baseline. These statewide values all incorporate an average value of vertical land motion corresponding to a negligible rate of 0.1 mm (0.0003 ft) per year uplift. The California Sea Level Scenarios track closely with global mean sea level (GMSL), with differences of only 2 to 3 inches between GMSL and the California Sea Level Scenarios in 2100. Evaluation of the Intermediate, Intermediate-High, and High scenarios (outlined in red below) is recommended to inform appropriate sea level rise planning and project decisions.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 |
| 2030 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 |
| 2040 | 0.4 | 0.5 | 0.6 | 0.7 | 0.8 |
| 2050 | 0.5 | 0.6 | 0.8 | 1.0 | 1.2 |
| 2060 | 0.6 | 0.8 | 1.1 | 1.5 | 2.0 |
| 2070 | 0.7 | 1.0 | 1.4 | 2.2 | 3.0 |
| 2080 | 0.8 | 1.2 | 1.8 | 3.0 | 4.1 |
| 2090 | 0.9 | 1.4 | 2.4 | 3.9 | 5.4 |
| 2100 | 1.0 | 1.6 | 3.1 | 4.9 | 6.6 |
| 2110 | 1.1 | 1.8 | 3.8 | 5.7 | 8.0 |
| 2120 | 1.1 | 2.0 | 4.5 | 6.4 | 9.1 |
| 2130 | 1.2 | 2.2 | 5.0 | 7.1 | 10.0 |
| 2140 | 1.3 | 2.4 | 5.6 | 7.7 | 11.0 |
| 2150 | 1.3 | 2.6 | 6.1 | 8.3 | 11.9 |



Key Takeaways

The California Sea Level Scenarios show greater certainty in the amount of sea level rise expected in the next 30 years than previous reports and demonstrate a narrow range across all possible emissions scenarios. Statewide, sea levels are most likely to rise 0.8 ft (Intermediate Scenario) by 2050.

In the mid-term (2050-2100), the range of possible sea level rise expands due to more uncertainty in projected future warming from different emissions pathways and certain physical processes (i.e. rapid ice sheet melt). By 2100, statewide averaged sea levels are expected to rise between 1.6 ft and 3.1 ft (Intermediate-Low to Intermediate Scenarios), although higher amounts are possible.

Over the long-term (towards 2100 and beyond), the range of sea level rise becomes increasingly large due to uncertainties associated with physical processes, such as earlier-than-expected ice sheet loss and resulting future sea level rise. Sea levels may rise from 2.6 ft to 11.9 ft (Intermediate-Low to High Scenarios) by 2150, and even higher amounts cannot be ruled out.

Vertical land motion is the primary driver of local variations in sea level rise across the state, driven by a combination of tectonics, sediment compaction, and groundwater and hydrocarbon withdrawal. Vertical land motion is incorporated into the sea level scenarios for each National Oceanic and Atmospheric Administration (NOAA) tide gauge and illuminates locations experiencing subsidence or uplift.

The pathway associated with the extreme sea level rise scenario (i.e. H++) from Rising Seas 2017 is higher than the best available science now supports. The key lines of evidence that resulted in the extreme sea level rise scenario (i.e. H++) from Rising Seas 2017 have been updated and are now reflected in the Intermediate-High and High Scenarios.

Today's coastal storms provide a glimpse into our future in which storm events will become more damaging and dangerous as climate change and sea level rise continue. Coastal storms under future sea level scenarios will cause accelerated cliff and bluff erosion, coastal flooding and beach loss, and mobilization of subsurface contaminants. Sea level rise will increase the exposure of communities, assets, services and culturally important areas to significant impacts from coastal storms.

Sea level rise will increase the frequency of coastal flooding events, which occur when sea level rise amplifies short-term elevated water levels associated with higher tides, large storms, El Niño events, or when large waves coincide with high tides. California communities need to be aware of and prepared for a likely rapid increase in the frequency of coastal flooding in the 2030s, even beyond the increases in coastal flood frequency already occurring as a result of extreme storms.

Groundwater rise poses a threat to below-ground infrastructure and freshwater aquifers under future Sea Level Scenarios. In areas with shallow unconfined groundwater, the water table will generally rise with sea level, depending on local geomorphology. Rising groundwater may mobilize subsurface contaminants in soils, expose underground infrastructure to corrosive saltwater, and put freshwater aquifers at risk of saltwater intrusion. The low-lying Sacramento-San Joaquin Delta, which supplies fresh water to two-thirds of the state's population and millions of acres of farmland, is particularly vulnerable to saltwater intrusion into freshwater aquifers.

BOX 2:**Guidance on Planning for Sea Level Rise Using the California Sea Level Scenarios**

This stepwise process recommends a precautionary approach for incorporating Sea Level Scenarios into planning and projects that includes adaptation pathways to phase actions over time. These steps complement other State guidance documents that also provide stepwise approaches to conducting analyses that inform sea level rise planning and decision making.

>> STEP 1: Identify the nearest tide gauge

The report's appendices provide a map of the 14 tide gauges in California for which localized Sea Level Scenarios are presented that incorporate the localized effects of vertical land motion.

>> STEP 2: Evaluate planning and/or project time horizon(s)

Determine how long a given planning effort or project is intended to function. If it is not possible to plan or adapt for the entire time horizon from the outset, a phased adaptation approach can be taken that provides earlier time horizons for interim adaptation steps.

>> STEP 3: Choose multiple Sea Level Scenarios for vulnerability assessment

It is recommended to evaluate the vulnerability of people, natural resources and infrastructure under the Intermediate, Intermediate-High, and High scenarios. Analysis of 100-year storm conditions under Sea Level Scenarios is also recommended, **with wave-driven processes and storm surge being the most important components to consider.**

>> STEP 4: Conduct vulnerability assessment

Conducting a vulnerability assessment begins with creating exposure maps of sea level rise-induced inundation and flooding, which

can also incorporate coastal erosion and groundwater rise. Once the physical extent of exposure is determined, a sensitivity analysis will provide information on the potential impacts of that exposure. The final step in a vulnerability assessment is for a community to determine its adaptive capacity to the determined impacts.

>> STEP 5: Explore adaptation options and feasibility

A collaborative process including affected communities, stakeholders, and relevant regulatory bodies should explore feasible adaptation options.

>> STEP 6: Select phased adaptation approach and/or implement project

Following an assessment of adaptation options, a specific project or adaptation pathway must be selected and implemented.

As the stepwise approach to applying Sea Level Scenarios is undertaken, general recommendations and principles to incorporate include: prioritization of social equity, environmental justice and the needs of underserved and vulnerable communities; protection of coastal habitats and public access; consideration of water-dependent infrastructure and uses; consideration of episodic increases in sea level rise caused by storms and other extreme events; coordination and collaboration with local, state and federal governments; consideration of local conditions; inclusion of adaptive capacity in design and planning; and assessment of risk and adaptation planning should be conducted at community and regional levels when possible.



1. Introduction

AUTHORS:
Ocean
Protection
Council
and Ocean
Science Trust

EARTH'S CLIMATE is changing, ice sheets and glaciers are melting, ocean water is warming and expanding, and sea levels are rising in response. The continuing and increasing emission of greenhouse gasses, particularly carbon dioxide, from the burning of fossil fuels (i.e., coal, oil, and natural gas) are the primary drivers of a warming planet. The more greenhouse gasses that we emit, the warmer the atmosphere and ocean become, and the higher and more rapidly sea level will rise in response.

The potential loss or damage to public infrastructure, health and safety, private developments, water supply, and natural habitats due to sea level rise is significant and urgent. Approximately 700,000 residents and \$250 billion in property across California could be exposed to the combination of storms and sea level rise-driven flooding during the 21st century.² Importantly, adverse impacts from sea level rise will have disproportionate impacts on vulnerable communities and may exacerbate existing environmental and systemic injustices. As we better understand the influence of climate change on the melting of glaciers and ice sheets, ocean warming, and their potential impact on future sea levels and associated coastal hazard risk, the better we can predict and estimate future sea level rise and inform adaptation planning.

As sea level rise accelerates, risks from other coastal processes are expected to increase. For example, rising sea levels will lead to more

2. Barnard et al., 2019.

episodic flooding and permanent inundation of low-lying areas that are exposed to high tides, and erode important habitats such as beaches, coastal wetlands, cliffs, and dunes. Higher sea levels will also contribute to rising groundwater tables that will bring saltwater intrusion into freshwater aquifers, increase corrosion of underground infrastructure, mobilize subsurface contaminants, and lead to surface pooling of water. In particular, the low-lying Sacramento-San Joaquin Delta supplies fresh water to two-thirds of the state's population and millions of acres of farmland and is vulnerable to groundwater aquifer saltwater intrusion.

Elevated sea levels can exacerbate the damaging effects of coastal storms. On January 5, 2023, wave heights reached 28 feet offshore from Monterey Bay, and arrived simultaneously with spring high tides. Strong onshore winds, which drive and add force to local storm waves and swells from distant storms, further elevated wave heights. Waves eroded the 20-30-foot-high bluffs along West Cliff Drive in Santa Cruz, destroying sections of the heavily used roadway and pedestrian path and endangering public safety. The co-occurring storm surge, large wave and high tide conditions also destroyed the historic wooden pier at Seacliff State Beach, flooded coastal streets and roads, and damaged both private development and public infrastructure along the northern Monterey Bay shoreline. Such extreme storms are anticipated to increase in frequency and intensity due to climate change.³ The cumulative impact of these storm impacts, combined with accelerating sea level rise, will result in even more significant damage along California's shoreline, including impacts to water supply and transportation infrastructure.

1.1. Existing State Policy Guidance and Sea Level Rise Planning

Over the past five years, sea level rise planning in California has been guided by the 2018 State of California Sea-Level Rise Guidance. In that guidance, OPC committed to updates approximately every five years to ensure that adaptation planning and projects are based on the best available science.

Since 2018, state agencies and departments have advanced sea level rise adaptation through the development, uptake, and implementation of multiple sea level rise policies, programs, and actions. For example, in 2020 the California Natural Resources Agency and California Environmental Protection Agency released a set of Sea Level Rise Principles with the purpose of aligning state planning, policy setting, project development, collaboration, and decision-making around sea level rise.⁴ These Principles were co-developed

3. Bromirski, 2023.

4. California Natural Resources Agency, 2020. http://www.opc.ca.gov/webmaster/_media_library/2020/05/State-SLR-Principles_FINAL_April-2020.pdf

with the State Sea Level Rise Collaborative, which is a group of 17 state agencies⁵ that meet quarterly to discuss coastal resilience issues at the state level, including emerging science, policy, and projects. Previously named the Sea Level Rise Leadership Team, the State Sea Level Rise Collaborative is also synonymous with the California Sea Level Rise State and Regional Support Collaborative, as referenced in Senate Bill 1 (Atkins, 2021). Facilitated by OPC, the Collaborative co-produced the State Agency Sea Level Rise Action Plan (Action Plan) for California in 2022, which provides a roadmap for coordinated and aligned state agency efforts to build resilience. The Collaborative provided an update to the Action Plan in February 2024.⁶

Since 2018, sea level rise planning efforts across the state have advanced in number, scale, and sophistication. Many coastal cities and counties have completed vulnerability assessments, which estimate the threat that sea level rise poses to public infrastructure, private homes, businesses, recreation areas, community centers, coastal habitats, and more.⁷ However, most of this planning has not yet considered the potential for impacts from rising groundwater or compound flooding. Sea level rise projections have been incorporated into local and regional planning and decision frameworks including Local Coastal Programs (LCPs), hazard mitigation plans, and the Delta Stewardship Council Delta Plan and Delta Adapts, among others.

In recent years, California has passed legislation aimed at advancing sea level rise adaptation planning throughout the state. Senate Bill 1 (SB 1, Atkins, 2021)⁸, the Sea Level Rise Mitigation and Adaptation Act, was signed into law in 2021. SB

1 directs OPC to administer grants to local and regional governments to plan for sea level rise and implement adaptation projects. With SB 1 funding, OPC established the SB 1 Sea Level Rise Adaptation Planning Grant Program (SB 1 Grant Program)⁹ that includes a technical assistance component, with the goal to provide funding for coastal communities to develop consistent sea level rise adaptation plans and projects to build resilience to sea level rise along the entire coast and San Francisco Bay. Additionally, Senate Bill 272 (SB 272, Laird, 2023)¹⁰ was signed into law in 2023. SB 272 requires that all coastal local governments develop sea level rise plans and prioritizes local governments with approved sea level rise plans for adaptation funding. These plans must be integrated into LCPs or San Francisco Bay shoreline resiliency plans by 2034. SB 272 also requires the California Coastal Commission (CCC), San Francisco Bay Conservation and Development Commission (BCDC), and OPC to establish minimum guidelines for sea level rise plans by 2024. Actions 4.1-4.3 in the State Agency Sea Level Rise Action Plan directly address and support these SB 272 requirements.

California continues to plan for the risks sea level rise poses to the state's Public Trust uses, which include commerce, navigation, fisheries, recreation, and conservation. State agencies have collaborated on guidance and recommendations for protecting Public Trust uses along the open coast and there are plans to similarly study resiliency strategies and create guidelines for Public Trust uses within the San Francisco Bay. The California State Lands Commission released a report in 2023, Shoreline Adaptation and the Public Trust: Protecting California's Public Trust

5. State Sea Level Rise Collaborative members: California Natural Resources Agency, California Environmental Protection Agency, San Francisco Bay Conservation and Development Commission, California Coastal Commission, California Energy Commission, California Department of Fish and Wildlife, Caltrans, Delta Stewardship Council, Department of Water Resources, Ocean Protection Council, Governor's Office of Planning and Research, Office of Emergency Services, State Coastal Conservancy, State Lands Commission, State Parks, State Water Resources Control Board, Strategic Growth Council, Department of Insurance.

6. Sea Level Rise Collaborative, 2024. <https://opc.ca.gov/wp-content/uploads/2024/02/SLR-Action-Plan-2024-Update-508.pdf>

7. Lester et al., 2023.

8. Senate Bill 1 https://leginfo.legislature.ca.gov/faces/billTextClient.xhtml?bill_id=202120220SB1

9. OPC SB 1 Funding Webpage <https://www.opc.ca.gov/sb-1-funding>

10. Senate Bill 272 https://leginfo.legislature.ca.gov/faces/billNavClient.xhtml?bill_id=202320240SB272

Resources from Sea Level Rise, which analyzes how sea level rise and adaptation strategies affect Public Trust resources and updates the state’s leasing practices related to shoreline protection structures and management of coastal state lands and Public Trust resources.¹¹ Additionally, in 2023 the CCC adopted the Public Trust Guiding Principles & Action Plan to describe how the Public Trust doctrine relates to sea level rise planning under the Coastal Act.¹² This report analyzes the policy, legal, and coordination issues raised by the challenges of protecting Public Trust lands, uses, and resources as sea levels rise.

1.2. Purpose & Intended Use

This report, and the accompanying data sets and links to tools and resources, serves as the 2024 update to the previous 2018 guidance. This report aims to: (1) synthesize and provide updated science of sea level rise and coastal hazards; and (2) provide practical and pragmatic guidance for applying this updated science across planning efforts and decision-making contexts.

This report provides guidance for a broad range of audiences including state agencies, tribes, regional and local governments, and climate adaptation planners. It is intended to foster coordinated and consistent statewide planning and decision-making based on science, and to enable the incorporation of sea level rise into the full suite of relevant sectors, policy decisions, adaptation plans, project designs, and investments.

The information contained in this report is intended to cover and be relevant for tidally influenced areas of the California shoreline; this includes all of California’s outer coast and parts of the Sacramento-San Joaquin River Delta. For more information on the specific geographic scope of the report, the spatial footprint of NOAA’s VDATUM tool can be used.¹³ VDATUM is

designed to convert spatial data among different vertical datums including tidal, orthometric, and ellipsoidal datums. Application of the Sea Level Scenarios provided here requires coupling to a vertical datum, and thus the ability to produce relevant datums for a given location can be viewed as a necessary requirement of applicability. The extent inland typically follows the reach of tidal influences along waterways that connect to the ocean. For California, this includes parts, but not all, of the Sacramento-San Joaquin River Delta, which has significant tidal effects and where marine and freshwater systems meet (see Box 3 for more information). For more specific information on the spatial extent covered, users are referred to the VDATUM tool and associated supporting information.

BOX 3:

Sacramento-San Joaquin Delta

The Sacramento-San Joaquin Delta stands out among California’s coastal regions due to its intricate network of waterways, islands, and marshlands, serving as a crucial nexus for water supply, agriculture, and biodiversity. Unlike typical coastal areas, the Delta’s hydrology—driven by both ocean conditions and inflows from five different river systems and a complex system of levees and channels, already under strain from subsidence and seismic activity — presents unique challenges for accurately assessing the impacts of climate change. The guidance provided in this document should be used in conjunction with an understanding of the Delta’s hydrological, geological, and environmental features when assessing climate change impacts to promote long-term sustainability.

11. California State Lands Commission, 2023. <https://slcprdwordpressstorage.blob.core.windows.net/wordpressdata/2023/12/Shoreline-Adaptation-Report.pdf>

12. California Coastal Commission, 2023. https://documents.coastal.ca.gov/assets/public-trust/Public%20Trust%20Guidance%20and%20Action%20Plan_Adopted.pdf

13. <https://vdatum.noaa.gov/>

1.3. How This Report was Developed

In June 2022, OPC and OST convened an interdisciplinary Sea Level Rise Science Task Force (hereafter referred to as the Task Force) to update the science foundation for state sea level rise policy. Task Force members contributed their technical expertise in meetings and discussions. OPC and OST collaborated to update the policy guidance as needed to incorporate this new science. OPC and OST consulted and gained input from state agency staff and local and regional planners, via multiple meetings and workshops, to align the updated science and guidance with their needs, interests, and opportunities for planning and preparing for sea level rise. The updated sea level rise science for California in Chapter 2.0 of this report was peer reviewed by three independent experts to ensure technical rigor and was revised in response to reviewer comments.

Feedback from the State Sea Level Rise Collaborative was incorporated prior to a public comment period. OPC conducted initial listening sessions and formal consultation with California Native American tribes to ensure that the guidance reflects tribal priorities and meets the needs of tribal governments and tribal communities. This final version reflects edits made in response to public comments received during a 45-day public comment period.

1.4. Updating the Science Foundation for Statewide Policy Guidance

The 2018 California Sea-Level Rise Guidance was based on a science synthesis, the *Rising Seas in California* report (hereafter referred to as *Rising Seas 2017*), developed by an expert panel and released the previous year. *Rising Seas 2017* provided sea level projections for specific locations along the California coastline, covering the time period from 2020 to 2150. Three different probabilistic projections tied to “Representative Concentration Pathways”, or RCPs, were provided alongside an extreme sea level rise scenario with unknown probability referred to as H++.

Since the release of *Rising Seas 2017*, the scientific community has made significant improvements in its ability to understand and project future sea level rise, offering a timely opportunity to update California’s sea level rise projections and align state guidance with this

new science. In particular, observational and modeling studies have provided more clarity on when and how much ice sheet and glacial melt will contribute to future sea level rise. This improved scientific understanding has been reflected in updated global probabilistic sea level rise projections provided by the Intergovernmental Panel on Climate Change 6th Assessment Report (IPCC AR6).¹⁴ In February 2022, a multi-agency federal group delivered the 2022 Federal Sea-Level Rise Technical Report¹⁵, which integrated this best available science from IPCC AR6 into five sea level scenarios that span the plausible range of sea level rise from 2020 to 2150 for all U.S. states and territories.

14. Masson-Delmotte et al., 2021.

15. Sweet et al., 2022.

BOX 4:**Representative Concentration Pathways (RCPs)**

Future greenhouse gas emissions and concentrations are difficult to predict and depend on future developments such as future population growth, economic growth, energy use, uptake of renewable energy, technological change, deforestation and land use. The climate-modeling community developed four RCPs that span a large range of future global warming scenarios. RCPs are space and time dependent trajectories of future greenhouse gas concentrations and different pollutants caused by different human activities. RCPs quantify future greenhouse gas concentrations and the radiative forcing (additional energy taken up by the Earth system), due to increases in greenhouse gas emissions.

Shared Socioeconomic Pathways (SSPs)

Developed more recently, the SSPs are a collection of narrative descriptions of alternative futures of socio-economic development in the absence of climate policy intervention. Five SSPs describe five different pathways that the world could take, drawing on data including population, economic growth, education, urbanization, and the rate of technological development. The SSPs are important inputs into the IPCC sixth assessment and are used to explore how societal choices will affect greenhouse gas emissions.

1.4.1 Framing potential sea level rise futures through scenarios

In Chapter 2.0 of this report, the Task Force adopted the scenarios framework in the 2022 Federal Sea-Level Rise Technical Report to provide Sea Level Scenarios for California, bringing consistency to state and federal sea level rise planning. These scenarios are derived from the sets of probabilistic projections developed in the IPCC AR6 and reflect the most up to date understanding of the physical drivers of sea level rise. As such, the information about the likelihood of meeting or exceeding a specific Sea Level Scenario is embedded in the scenarios themselves (e.g., the High Scenario is less likely than the Intermediate Scenario, as described in more detail in Chapter 2.0).

The Sea Level Scenarios for California span the plausible range of future sea level rise under all of the possible emissions and global development futures, or SSPs, defined by the IPCC AR6. While Rising Seas 2017 provided probabilistic projections under three IPCC emissions scenarios termed RCPs, the Scenarios in this report span all SSPs and enable users to consider sea level rise without first selecting a single emissions future on which to base planning.


1.5. Translating Science into Policy Guidance

California has a strong existing foundation for developing science-based, pragmatic policy guidance, even in the face of uncertainties about the future. Chapter 4.0 of this report applies the newest science on sea level rise and other coastal hazards to update policy guidance, as appropriate. **These updates do not represent new or significantly changed state policy direction.** Consistent with the 2018 California Sea-Level Rise Guidance, this 2024 update recommends a stepwise process for incorporating Sea Level Scenarios into planning and decisions that is still precautionary in nature. This report describes considerations for adaptation and planning that match the advancing sophistication of adaptation planning across the state and upholds California's values and priorities that are as relevant today as five years ago when the 2018 California Sea-Level Rise Guidance was released.

Rising Seas 2017 provided a rigorous treatment of scientific uncertainties associated with the physical processes of sea level rise by providing probabilistic projections. The 2018 California Sea-Level Rise Guidance then created decision-ready Sea Level Scenarios by selecting certain projections and assigning levels of risk aversion associated with each (see Table 1 in the 2018 California Sea-Level Rise Guidance). Like the 2018 sea level projections, the California Sea Level Scenarios in Chapter 2.0 incorporate a risk-based framing by addressing uncertainties

in the physical processes causing sea level rise, and in the ability to model and project future timing and magnitudes of sea level rise. Chapter 2.0 differs from the 2018 California Sea-Level Rise Guidance in that it provides Sea Level Scenarios spanning a range of emissions pathways, rather than several sets of probabilistic projections each linked to a different emissions pathway (RCP). Like the 2018 California Sea-Level Rise Guidance, the stepwise process for incorporating sea level rise into planning and decisions still includes consideration of other important elements of risk, qualitative and quantitative social and economic impacts, and adaptive capacity.

Chapter 3.0 of this report describes how sea level rise intersects with multiple and often compounding hazards including coastal and precipitation-based flooding, groundwater rise, and shoreline change. Application and use of Sea Level Scenarios and the accompanying policy guidance is not a single step but should be invoked at multiple stages of climate adaptation planning and implementation. To that end, several data visualization and decision-support tools have been developed, and continue to be refined, to support local, regional, and statewide adaptation planning, which are summarized in Appendix 4. OPC, working with local, tribal, state and federal partners, will continue to support and advance uptake and use of this policy guidance in the full breadth of adaptation planning and implementation that intersects with sea level rise.



BOX 5:**Causes of Sea Level Rise in California**

Sea level rise along the coast of California is driven by a combination of processes. These processes operate on different spatial scales – from global to local – and affect the California coastline in different ways.

Global Sea Level Rise

The primary causes of sea level rise on global scales are thermal expansion due to ocean warming and the input of freshwater from melting ice sheets and glaciers on land. Both of these processes result from ongoing warming of the planet due to greenhouse gas emissions from human activities. Tide gauge measurements show roughly 5 inches of GMSL rise during the 20th century.¹⁶ Since 1993, satellite altimeters have provided continuous near-global measurements of sea level, showing an additional 4 inches of GMSL rise just in the past 30 years.¹⁷ From these observations, it is clear that sea level rise is accelerating,¹⁸ and the current rate of GMSL rise (1.7 inches/decade) is triple the 20th century rate.

Regional Relative Sea Level Rise

Sea level rise is not uniform across the globe. Relative sea level rise (the rise of seas relative to land) at any specific location is driven by a combination of the global processes described above plus three primary local and regional processes:

1. Sterodynamic Sea Level Change:

Sterodynamic sea level change describes the combined effect of steric (temperature and salinity) changes and changes in ocean dynamics (i.e. winds, currents). Sea level rise caused by large-scale persistent changes in ocean dynamics is not currently expected to be consequential for the California coast.¹⁹ During the satellite altimeter era (1993-present), the combination of shorter-term El Niño-Southern Oscillation (ENSO, or El Niño) and decadal variability led to a

dramatic shift in the rates of sea level rise during the satellite record, which have now evened out over the full 30-year record. These shifts will continue to happen in the future, but persistent sterodynamic sea level rise along the California coast is expected to be primarily driven by ongoing thermal expansion of the ocean and, therefore, will track closely with the global average into the future.²⁰

2. Gravitational, Rotational, and Deformational:

The impact of ice-mass loss is expected to be larger for the California coast than the global average.²¹ As an ice sheet loses mass to the ocean, its gravitational pull on the surrounding ocean is reduced. In the vicinity of the ice sheet, the reduced gravitational pull on the ocean causes the sea level to decrease, but as distance from the ice sheet increases, the change in local relative sea level becomes greater than the global average²² (see *Rising Seas 2017* or Hamlington et al., 2020 for detailed explanation). As a result, California is heavily impacted by ice loss from the West Antarctic Ice Sheet, and for every foot of global sea level rise caused by ice loss in West Antarctica, California sea level will rise about 1.25 feet. The Greenland Ice Sheet, on the other hand, contributes about 0.75 feet of sea level rise to California for every foot of global sea level rise. Due to its larger-scale regional impact, GRD effects do not lead to large differences in sea level rise between locations along the California coast.

16. Frederikse et al., 2020.

17. Willis, Hamlington, Fournier, 2023.


18. Dangendorf et al., 2019; Nerem et al., 2018.

19. Fox-Kemper et al., 2021.

20. Hamlington et al., 2021.

21. Griggs et al., 2017.

22. Mitrovica et al., 2011.

BOX 5 CONT.

3. Vertical Land Motion: Vertical land motion is the result of a combination of processes acting together on different temporal and spatial scales. On the largest scales, Earth's surface is still rising and falling after the retreat of large ice sheets that covered the Northern Hemisphere about 18,000 years ago, a process referred to as glacial isostatic adjustment.²³ Given the distance of California from these past ice sheets, the impact of these land adjustments is small, but detectable. On a local scale, tectonics, sediment compaction, and groundwater and hydrocarbon withdrawal play a role in vertical land motion in California. Records of vertical land motion observation are short, and a complete understanding of vertical land motion at any particular location generally requires considerations of local land-use alongside available measurements. When compared to sterodynamic and gravitational, rotational, and deformation-related changes in sea level rise along the California Coast, the impact of vertical land motion on relative sea level rise is much more localized and can lead to significant differences from one location to the next.

23. Peltier et al., 2004.

2. California Sea Level Scenarios

LEAD AUTHOR: Dr. Benjamin Hamlington

SUPPORTING AUTHORS: Dr. Felix Landerer, Ocean Science Trust

CONSULTATION PROVIDED BY: Dr. Susheel Adusumilli, Dr. Patrick Barnard, Dr. Dan Cayan, Laura Engeman, Dr. Gary Griggs, Dr. Kristina Hill, Dr. Phil Thompson

BOX 6:

California Sea Level Scenarios

- **Since Rising Seas 2017**, the scientific community has made significant improvements in its ability to understand and project future sea level rise and this best available science was incorporated, in this report, into a set of five California Sea Level Scenarios from 2020 to 2150.
- **The California Sea Level Scenarios** show greater certainty in the amount of sea level rise expected in the next 30 years than previous reports and demonstrate a narrow range across all possible emissions scenarios. Statewide, sea levels are most likely to rise 0.8 ft (Intermediate Scenario) by 2050.
- **In the mid-term** (2050-2100), the range of possible sea level rise expands due to more uncertainty in projected future warming from different emissions pathways and certain physical processes (i.e. rapid ice sheet melt). By 2100, statewide averaged sea levels are most likely to rise between 1.6 ft (Intermediate-Low Scenario) and 3.1 ft (Intermediate Scenario), although higher amounts are possible.
- **Over the long-term** (towards 2100 and beyond), the range of sea level rise becomes increasingly large due to uncertainties associated with physical processes, such as earlier-than-expected ice sheet loss and resulting future sea level rise. Sea levels may rise from 2.6 ft (Intermediate-Low Scenario) to 11.9 ft (High Scenario) by 2150, and even higher amounts are possible.
- **Vertical land motion** is the primary driver of local variations in sea level rise across the state, driven by a combination of tectonics, sediment compaction, and groundwater and hydrocarbon withdrawal. Vertical land motion is incorporated into the Sea Level Scenarios for each National Oceanic and Atmospheric Administration (NOAA) tide gauge and illuminates locations experiencing subsidence or uplift.
- **The key lines of evidence** that resulted in the extreme sea level rise scenario (i.e. H++) from Rising Seas 2017 have been updated and are now reflected in the Intermediate-High and High Scenarios.

IN AUGUST 2021, the Working Group I contribution to the IPCC AR6 was released.²⁴ This contribution provides the most up-to-date physical understanding of the climate system and climate change, bringing together the latest advances in climate science. Chapter 9 of that contribution provides the latest scientific understanding of the physical processes underlying global and regional changes in the ocean, cryosphere and sea level²⁵ and forms the scientific foundation for developing California Sea Level Scenarios.

The 2022 Federal Sea Level Rise Technical Report includes a set of five sea level scenarios for the U.S., providing a range of plausible changes through 2150. The scenarios were developed from the suite of modeled projections in the IPCC AR6 that include new advancements in the understanding of when and how various global and regional processes may occur (e.g., ocean dynamics, glacier and ice sheet melt, mass redistribution). The five scenarios (Low, Intermediate-Low, Intermediate, Intermediate-High, and High) correspond to average global sea level rise magnitudes in the year 2100. Here, the same framework and approach is adopted for consistency, and the scenarios are regionalized to develop California-specific scenarios.

2.1. Updated Scientific Understanding

Since *Rising Seas 2017* was developed, scientific understanding of both present and future sea level has evolved. Recent observations of sea level change have advanced the scientific understanding of present sea level rise, and advances in the projection of future sea level rise reflect a deeper understanding of possible high-end estimates of future sea level rise.

2.1.1. Observed sea level change

The rate of sea level rise along California's shoreline during the satellite altimeter era (1993-present) has been less than the global average for much of that record²⁶ due primarily to the influence of natural variability temporarily obscuring the background, climate-driven rate. The combination of El Niño and decadal variability associated with the Pacific Decadal Oscillation has led to dramatic

shifts in the rate of sea level rise across the 30 years of the satellite record, wherein sea level rise was essentially absent during the first half and was substantial during the second half of the record (Figure 2.1, top). These shifts will likely continue in the future, but longer tide gauge records together with recent observations suggest that sea level rise along the California coast should resemble the global average.²⁷ The time series of average California sea level change from both tide gauges and satellite altimetry is shown in Figure 2.1 (bottom). Over the complete record from 1993 to 2023, the rate of sea level rise for California, on average, is 0.9 inches/decade.

2.1.2. Advances in projecting future sea level rise

The current consensus understanding on sea level rise was described in the IPCC AR6, which used this knowledge to create new projections of future sea level rise. The updated IPCC AR6 sea level projections are formed by

24. Masson-Delmotte et al., 2021.

25. Fox-Kemper et al., 2021.

26. Bromirski et al., 2011; Thompson et al., 2014; Moon et al., 2015; Hamlington et al., 2021.

27. Hamlington et al., 2021.

integrating different projections of individual processes that cause sea level change within a consistent framework.²⁸ The latest generation of global climate models from the Coupled Model Intercomparison Program’s sixth phase (CMIP6) are used to account for the ocean

dynamic regional sea level rise and similar methods are used for assessing vertical land motion contributions as in past reports. For the coast of California, the updates do not lead to significant changes for these contributions relative to past reports.




FIGURE 2.1. Top: The rate of sea level rise, in inches per decade, for California estimated from both tide gauges and satellite altimetry over three different time periods: 1993-2008, 2008-2023, and 1993-2023. **Bottom:** Sea level variations in feet averaged along the California coastline from both tide gauges and altimetry.

By comparison, the IPCC AR6 reflects a step forward in scientific understanding of the ice sheet contributions to sea level rise, and in how this understanding should be incorporated into projections. This has altered projections of sea level rise relative to past reports (e.g., Rising Seas 2017). In addition to being important sources of future sea level rise, projections of future ice sheet change represent the largest sources of uncertainty in estimating sea level rise towards the end of this century and beyond, even with this new understanding. By comparison, the IPCC AR6 reflects a step forward in scientific understanding of the ice sheet contributions to sea level rise, and in how this understanding should be incorporated into projections. This has altered projections

of sea level rise relative to past reports (e.g., Rising Seas 2017). In addition to being important sources of future sea level rise, projections of future ice sheet change represent the largest sources of uncertainty in estimating sea level rise towards the end of this century and beyond, even with this new understanding.

Within the AR6 framework, a set of SSP scenarios that project global socioeconomic changes to 2100 are used to drive models of the physical processes causing sea level rise to

28. See Table 9.7; Fox-Kemper et al., 2021.

generate sea level projections. Five SSP-based projections included only physical processes in which there is at least medium confidence in the current scientific understanding, and two additional scenarios (one low emissions, one high emissions) included ice sheet processes in which there is currently low confidence among scientists (see Box 7 for a description of ‘confidence’). These low confidence processes include earlier-than-projected ice-shelf disintegration in Antarctica, abrupt and widespread onset of marine ice-sheet instability and/or marine ice-cliff instability in Antarctica, and faster-than-projected changes in surface-mass balance on Greenland. Low confidence as applied to these processes reflects an incomplete scientific understanding of the physics underlying these processes, and associated gaps in representing these processes in the current generation of models that are used for projections. This leads to low confidence in the ability to quantify the sea level rise that will result if any of these processes were to be triggered. As a result of the low confidence in these processes, the two SSP scenarios in which they appear are considered of unknown likelihood.

2.1.3. Timing of high-end sea level rise

Although the IPCC AR6 assigned low confidence in the role these processes will play in the future, they do play an important role in determining the full range of possible sea level rise at any time in the future. An important change in the IPCC AR6 projections resulted from an update in the understanding of when these low confidence processes could come into play. The physical processes that could lead to much higher increases in sea level are now viewed by the scientific community as less plausible in the coming decades before potentially becoming a factor towards the end of the 21st century and beyond.²⁹ Additionally, at lower future warming levels (less than 2°C by 2100), significant contributions from these processes are not expected until beyond 2100, if at all.

As a result, when compared to past reports such as Rising Seas 2017, there is less acceleration of sea level before 2050, and the possibility of greater acceleration only towards the end of the 21st century and beyond. This has two primary implications when compared to the projections in the 2018 California Sea-Level Rise Guidance. First, there is now a narrower range of plausible sea level rise prior to 2050. Second, the lines of evidence used to construct the H++ scenario in Rising Seas 2017 have been updated, which leads to a shift in the timing of possible high-end contributions to sea level rise from the ice sheets. The previous description of future sea level rise described by H++, is not physically plausible as it incorporates too much sea level rise in the near-term and a consequent ongoing high rate of sea level rise throughout the rest of the 21st century.

29. DeConto et al., 2021; DeConto & Pollard, 2016.

BOX 7:**Confidence**

Confidence, in IPCC terms, measures a combination of available evidence and agreement among scientists. Evidence assesses the amount, quality and consistency of lines of evidence agreeing with a conclusion, while agreement evaluates the breadth of support for conclusions among experts. The IPCC AR6 uses a range of qualifiers to express this assessment, and therefore, confidence: Very Low, Low, Medium, High, and Very High.

In this report, two types of confidence are used:

- 1. Medium Confidence:** used to denote moderate agreement among experts on the model treatment of key processes (e.g. those used in the IPCC AR6 SSPs) and moderate lines of evidence supporting model outputs.
- 2. Low Confidence:** used to denote a low level of agreement on how models represent key processes (e.g. partial, rapid ice sheet disintegration) and limited evidence supporting model outputs.

2.2. Steps to Build California Sea Level Scenarios

Five Sea Level Scenarios have been developed for the California coast using the following process, also depicted in schematic form in Figure 2.2. This process is consistent with the approach in the 2022 Federal Sea Level Rise Technical Report, but is regionalized to produce California-specific scenarios. A description of their formation is provided here, and additional detail that could assist in interpretation is found in Appendix 4:

1. A front-end assessment of the projections and science contained in IPCC AR6 is used to determine a plausible (see Box 8 for definition) range of global mean sea level rise of between 0.3 and 2.0 m in 2100. Further support for the selection of the plausible range is provided in Figure 2.4 through analysis of the ranges of the AR6 scenarios at different time horizons.
2. The IPCC AR6 created medium confidence sea level projections for multiple SSPs.

In addition, low confidence projections allowing for potential contributions from rapid ice-sheet loss processes with unknown likelihood were also created for the SSP1-2.6 and SSP5-8.5 scenarios. For each SSP, an ensemble of thousands of “samples” of the trajectory of sea level rise from 2020 to 2150 was produced. These samples cover a range of possibilities, and the associated distribution of these samples for each SSP is shown graphically on the left of Figure 2.2.

3. Five GMSL targets or “gates” in 2100 that span the plausible range from 0.3 to 2.0 m are defined. The full set of samples is filtered to find the ones that fit through each of these five gates +/- 2 cm. As shown in Figure 2.2, the SSPs with the lowest emissions primarily build the lowest sea level scenarios. The SSPs with the highest emissions plus the scenario that takes into account rapid ice sheet loss primarily build the higher scenarios. These GMSL targets used to define the California Sea Level Scenarios are consistent with the national report.



4. Finally, five time-varying Sea Level Scenarios of GMSL and associated local relative sea level rise are constructed by collecting all the samples from across SSPs that fit through the defined gates. Logically, the Intermediate Low and Intermediate scenarios will include the highest number of samples, while the other sea level scenarios will have comparatively less. The individual samples of sea level projections provide important contextual information for each of the Sea Level Scenarios, carrying with them a range of warming levels and emissions pathways. While it is not possible to directly assign probabilities to each of the sea level scenarios, additional assumptions about the future (e.g. warming level) allow for the assignment of a probability of exceeding a particular Sea Level Scenario in that assumed future (see Chapter 2.4).

The contributions from steric sea level rise and from ice mass loss are similar across all of California, varying by less than 1.2 inches from the southern to northern extents at all time periods and all scenarios prior to 2100; vertical land motion is the primary driver of variations among different locations across the state. A single average value of vertical land motion (corresponding to a negligible

rate of 0.10 mm/year or 0.04 inches/decade uplift) is used for this chapter where statewide values of the Sea Level Scenarios are provided. Sea Level Scenarios for individual NOAA tide gauge locations that incorporate site-specific vertical land motion projections can be found in Appendix 2.

The scenarios show the rise in mean relative sea level over time and represent only the relative sea level rise happening on long timescales. Shorter period sea level change associated with El Niño, tides, storm conditions, or other natural ocean variability is not included in the scenarios themselves and does not contribute to the likely range of the scenarios. Including additional sea level rise on top of these scenarios (i.e., a buffer) to account for this natural variability is not recommended. Adding a buffer on top of the scenarios to generate new target values can lead to misleading interpretations and conclusions and could even lead to double-counting of natural sea level variability when subsequently projecting flood frequency and severity. Rather, the contributions from these temporary fluctuations can be considered related to coastal hazards that contribute to sea levels or when end users are required to use an online tool with limited sea level rise value options (see Chapter 3.0 for more detail).




FIGURE 2.2. Schematic showing that the construction of the sea level scenarios is based on SSPs, which inform a range of plausible future sea level rise. The full range of plausible global sea level rise in 2100 is then divided into sea level scenarios ranging from low to high. The final scenarios cover the time period from 2020 to 2150.

2.2.1 Evolving understanding of ice sheet instability

Some studies published after the release of the IPCC AR6, and garnering significant media attention, suggest that the onset of rapid ice sheet loss may happen in the coming decades.³⁰ Even if this does occur, there is large uncertainty as to the total extent of the ice sheet loss that would occur as a result and – more importantly – there is large uncertainty as to how quickly sea levels would rise in response, with most timelines beyond the end of the 21st century. Other recent studies have further assessed these low confidence processes to obtain a practical high-end estimate of future sea level, obtaining estimates within the range of the IPCC AR6 projections.³¹ Thus, the scenarios developed here for California represent the best available scientific understanding as described both in the IPCC AR6 and in the studies released subsequently.

30. E.g., Box et al., 2022; Stokes et al., 2022.

31. van de Wal, 2022.

2.2.2. Measuring vertical land motion

Satellite radar observations from 2014 to present have recorded land movement ranging from 2 inches of sinking or lowering to 2 inches of uplift, with some locations exceeding this typical range.³² In this chapter of the report, a statewide average rate of vertical land motion was estimated using a statistical model that incorporated observational data from individual tide gauges. Across all locations, the long-term rate of past vertical land motion is assumed to persist into the future, and future relative sea level rise is adjusted accordingly. This average rate of vertical land motion is near zero (0.1mm or 0.003ft uplift per year). Furthermore, the effect of vertical land motion is considered equal across each of the five California sea level scenarios; there is no significant difference between the vertical land motion contribution for the Low and High scenarios (Note: apparent differences in the vertical land motion contributions across scenarios are due to a combination of rounding and the framework used for combining process contributions and generating the projections).

Beyond the statewide averages used in this chapter, there are local variations in the rate of vertical land motion that must be factored in. Scenarios for individual NOAA tide gauge locations that include the local estimates of vertical land motion produced by the IPCC AR6 and included in the 2022 Federal Sea Level Rise Technical Report are provided in Appendix 2. Alternatively, where very localized global navigation satellite systems (GNSS or GPS) data or information from satellite-based approaches is available allowing more resolved estimates of vertical land motion, these can be added to the statewide average scenario values provided in this chapter. Additional information for doing this is provided in Appendix 2.

2.3. Sea Level Scenarios 2020-2150

Five Sea Level Scenarios are constructed and presented for California. Each scenario is defined and labeled according to the target value of global mean sea level (GMSL) rise in 2100:

- **Low** (0.3m or 1.0ft by 2100)
- **Intermediate-Low** (0.5m or 1.6ft by 2100)
- **Intermediate** (1.0m or 3.3ft by 2100)
- **Intermediate-High** (1.5m or 4.9ft by 2100)
- **High** (2.0m or 6.6ft by 2100)

32. Govorcin et al., 2024

TABLE 2.1. (A) Median values (i.e., 50th percentile) for Sea Level Scenarios for California, in feet, relative to a 2000 baseline. These statewide values all incorporate an average value of vertical land motion corresponding to a negligible rate of 0.1 mm (0.0003 ft) per year uplift.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 |
| 2030 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 |
| 2040 | 0.4 | 0.5 | 0.6 | 0.7 | 0.8 |
| 2050 | 0.5 | 0.6 | 0.8 | 1.0 | 1.2 |
| 2060 | 0.6 | 0.8 | 1.1 | 1.5 | 2.0 |
| 2070 | 0.7 | 1.0 | 1.4 | 2.2 | 3.0 |
| 2080 | 0.8 | 1.2 | 1.8 | 3.0 | 4.1 |
| 2090 | 0.9 | 1.4 | 2.4 | 3.9 | 5.4 |
| 2100 | 1.0 | 1.6 | 3.1 | 4.9 | 6.6 |
| 2110 | 1.1 | 1.8 | 3.8 | 5.7 | 8.0 |
| 2120 | 1.1 | 2.0 | 4.5 | 6.4 | 9.1 |
| 2130 | 1.2 | 2.2 | 5.0 | 7.1 | 10.0 |
| 2140 | 1.3 | 2.4 | 5.6 | 7.7 | 11.0 |
| 2150 | 1.3 | 2.6 | 6.1 | 8.3 | 11.9 |

(B) Rates of sea level rise (inches/year) for California in 2050 and 2100

| SCENARIO | RATE IN 2050 (inches/year) | RATE IN 2100 (inches/year) |
|-------------------|-------------------------------|-------------------------------|
| Low | 0.1 | 0.1 |
| Intermediate-Low | 0.2 | 0.2 |
| Intermediate | 0.3 | 0.8 |
| Intermediate-High | 0.5 | 1.1 |
| High | 0.7 | 1.3 |




FIGURE 2.3. Sea Level Scenarios from 2020 to 2150, in feet, with a baseline of 2000. For comparison, the H++ from the 2018 California Sea-Level Guidance is included illustrating that this scenario is above scientifically plausible sea level rise for all dates.

The sea level rise values associated with five Sea Level Scenarios for California are shown in Table 2.1 and Figure 2.3. The key features of these five scenarios for California are noted here and discussed in more detail in the subchapters that follow:

- Through 2100, the scenarios for California track closely with global mean sea level (GMSL), with differences of only 2 to 3 inches between GMSL and the California Sea Level Scenarios in 2100.
- Taken together, the median values of the Sea Level Scenarios capture the plausible range of sea level rise for all time periods prior to 2100 (see Box 8 for description of plausible).
- Beyond 2100, the range of plausible sea level rise increases significantly and extends beyond that captured by the Low to High Scenarios, as the potential for low confidence processes to contribute to sea level increases.
- The rate of sea level rise in 2050 and 2100 associated with each of these scenarios is also shown in Table 2.1(B). To reach the higher scenarios by 2050, the rate of sea level rise along the California coast would have to increase dramatically from the rate of ~0.1 inches/year over the past 30 years. In 2100, the implied rate of sea level rise is greater than 1 inch/year for the higher-end scenarios.

BOX 8:**Understanding Sea Level Scenario likelihoods**

The terms “most likely” and “plausible” have been used in recent assessment reports and are again used here. To date, a definition for these terms and how they have been applied to assess likelihoods of sea level scenarios has not been provided. To assist in their interpretation and application, the terms are defined here.

Most Likely

The phrase “most likely” is used when a scientific assessment of multiple lines of evidence collectively and consistently points towards a single sea level scenario or range across multiple sea level scenarios.

These lines of evidence include: a) the projected global surface temperature based on current emissions policy and commitments; b) the current trajectory of globally averaged sea level rise; c) the current trajectory of regionally averaged sea level rise; d) the range across the five sea level scenarios; e) the likely ranges covered by the medium confidence projections in the IPCC AR6. Based on these lines of evidence, the most likely sea level rise will be narrow in the near-term (2050), and they support the selection of a single sea-level scenario. Towards the end of the 21st century, the criteria indicate that the most likely sea level rise is best represented by the range across two sea level scenarios. Beyond 2100, based on current scientific understanding, lack of suitable lines of evidence for support, the possible contribution of low confidence processes, and general level of deep uncertainty, an assessment of the most likely range is not advised. Where the term expected is used, this refers to the amount or range of sea level rise that has been evaluated to be most likely.

Plausible

The plausible range of sea-level rise is the credible and reasonable range of future sea level rise supported by published, peer-reviewed publications and the consensus assessment of the IPCC AR6.

Plausible does not mean “possible”, which instead has particular meaning for the evaluation of the upper and lower ends of the plausible range. The Sea Level Scenarios are intended to have similar plausibility at the upper and lower ends of the full range (i.e., the Low and High scenarios). However, there is still disagreement within the scientific community about the plausible high-end estimate of sea-level rise between now and 2150. The upper-end of the plausible range is thus defined such that it is supported by scientific consensus and areas of overlap in published studies to the greatest degree possible; given ongoing scientific research, this may be further refined in future assessments.

2.3.1 Comparison of California Scenarios, IPCC AR6 Sea Level Projections, and California 2018 Guidance Scenarios

The goal in defining Sea Level Scenarios for California is for the five scenarios to span the full plausible range of sea level rise at any time from 2020 to 2150 (see the definition of plausible above). As opposed to constructing a projection around a particular emissions pathway, the scenarios specify a targeted amount of global mean sea level rise at a time (2100) in the future. The trajectory for getting to that target value relies on the same science and projection framework from the IPCC AR6. In other words, while the values of the Sea Level Scenarios are fixed to span the plausible range of GMSL rise in 2100, the trajectory of the Sea Level Scenarios before and after that time is set by the underlying AR6 projections.

For example, in 2050 and 2100, the median values of the five California Sea Level Scenarios encompass the 17th to 83rd interval from the AR6 projections (Figure 2.4 A and B). In 2150, however, the range expands due to the potential contributions of the low confidence processes. In particular, the upper end (shown in Figure 2.4 as 83rd percentile) of the range for the AR6 SSP5-8.5 Low Confidence projection increases significantly. After 2100, the Sea Level Scenarios do have an associated range due to the spread of pathways after reaching their target values in 2100. The upper end (83rd percentile) of the range for the High Scenario encompasses the 83rd percentile of the SSP5-8.5 Low Confidence scenario. Rather than indicating a flaw in the Sea Level Scenario framing, this exemplifies the increasing disagreement among scientists regarding how sea level rise and the important physical processes may evolve further in the future under continuing warming.

Due to the rapid near-term increase in sea level rise that is required, the Extreme Risk Aversion scenario from the 2018 California Sea-Level Rise Guidance, based on H++, is no longer physically realistic. However, there is consistency between the other risk aversion scenarios from the 2018 California Sea-Level Rise Guidance and those presented here. For example, across all three time periods, the Low Risk Aversion scenario closely tracks the range covered by Low and Intermediate Scenarios in this report. The Medium-to-High Risk Aversion scenarios correspond closely to the Intermediate-High and High Scenarios in 2100 and 2150, although are substantially higher in 2050. This difference reflects the narrower and lower range across the more recent IPCC AR6 projections until 2050.




FIGURE 2.4. Comparison between the sea level scenarios in this report (blue), the sea level projections from the IPCC AR6 (purple),³³ and the values from the 2018 California Sea-Level Rise Guidance (red) in 2050 (A), 2100 (B) and 2150 (C). The horizontal dashed lines indicate the range covered between the median values of the Low and High Scenarios. The vertical lines extending from the median values represent the likely ranges (17th-83rd percentiles) for each projection. Note, the sea level scenarios are defined based on the plausible range in 2100, and the values in 2050 and 2150 are determined by the pathways and time evolutions of the AR6 projections.

33. Fox-Kemper et al., 2021.

2.3.2. Sea level scenario changes through time

NEAR-TERM (2020-2050)

The California Sea Level Scenarios in this report show much greater certainty in the amount of sea level rise expected in the next 30 years compared to the previous report. This demonstrates that there is little difference in the amount of sea level rise expected across all foreseeable emissions pathways over the next three decades. In the 2018 California Sea-Level Rise Guidance, the risk aversion scenarios covered a range of 1.1 to 2.7 feet in 2050 relative to 2000. As a result of the updated science in the IPCC AR6, the Low and High Scenarios provide bounds in 2050 of 0.5 ft and 1.2 ft, a range of only 8 inches. This narrowing is driven primarily by a reduction in the upper-end scenario, and sea level rise of 2.7 feet by 2050 is now considered physically unrealistic. In other words, the increase in the rate of sea level rise that is needed to reach this target is not possible based on the current understanding of the processes driving sea level rise.

Furthermore, the 2050 Sea Level Scenarios can be informed by the trajectory of current observed sea level rise for California. In the 2022 Federal Sea Level Rise Technical Report, the sea level rise trend derived from observations over a minimum of 50 years, was extrapolated as an additional line of evidence during this near-term time period. Figure 2.5 shows an observation extrapolation using the tide gauge data from 1970 to 2022 to estimate a rate and acceleration. To the extent possible, the influence of natural variability, like that associated with El Niño, is removed to provide a more direct comparison to the Scenarios. The observation extrapolation tracks on top of the Intermediate Scenario, and the associated range lies between the Intermediate-Low and Intermediate-High Scenarios. In short, it is reasonable to view the Intermediate Scenario as most representative of the sea level rise that is most likely to occur between now and 2050, and small deviations of 4 inches either side of this scenario can represent the full plausible range.




FIGURE 2.5. *Sea Level Scenarios for California, in feet, from 2020 to 2050 relative to a baseline of 2000. The past statewide average tide gauge record is used to assess the trajectory of near-term sea level rise and create an observation extrapolation for comparison to the scenarios. The shaded region, bounded by the red dotted line, represents the likely range for the observation-based extrapolation.³⁴*

MID-TERM (2050-2100)

Beyond the middle of this century, the differences between the Sea Level Scenarios become increasingly large and more closely associated with both differences in potential future greenhouse gas emissions and possible contributions from low confidence ice sheet processes. The Low and High Scenarios have median values of 1.0 ft and 6.6 ft, respectively, in 2100.

The value of the High Scenario is substantially lower than the H++ scenario in 2100 (10.2 ft), and the High Scenario is itself very high relative to the IPCC AR6 medium confidence

projections, which include all AR6 projections except the AR6 SSP5-8.5 Low Confidence projections (Figure 2.4B). The H++ value in 2100 would not be reached even when considering the high end (95th percentile) of the AR6 SSP5-8.5 Low Confidence scenario (7.5 ft in 2100).

Extending the observational extrapolation in Figure 2.5 to 2100 would give an increase of 2.5 ft relative to 2000, which would fall in the middle of the Intermediate-Low and Intermediate Scenarios, but with a much larger likely range than in 2050.

34. Nerem et al., 2022; Hamlington et al., 2022.

LONG-TERM (2100-2150)

Beyond 2100, the range across the five Sea Level Scenarios increases significantly, going from 1.4 ft to 11.9 ft between the Low and High Scenarios. The upper end of the range of the AR6 SSP5-8.5 Low Confidence scenario expands to almost 18 ft. This reflects the rapid acceleration in sea level rise that may occur should the low confidence processes become an important factor. Unlike the other time periods, the median values of the Sea Level Scenarios do not fully encompass the 17th-83rd percentiles of the AR6 projections (Figure 2.4C). The 83rd percentile of the High Scenario is required to encompass the 83rd percentile of the SSP5-8.5 Low Confidence scenario (Figure 2.4C), although it should be noted that plausibility in this case is supported by a limited set of scientific studies and is a result of the SSP5-8.5 Low Confidence scenario alone.

In a future where the low confidence processes are not considered to be a factor, the range between the Intermediate-Low and Intermediate Scenarios represents the plausible sea level rise. The Intermediate through High Scenarios then represent the range of possibilities should the low confidence processes come into play. Towards 2100 and beyond, ambiguity arising from these low confidence processes and socioeconomic factors plays a major role and drives the wide range across scenarios.³⁵ Ambiguity arising from the low confidence ice-sheet processes is not just in regards to *what extent* and *when* these processes will develop, but also in how they will contribute to sea level rise should they develop. This should be considered when applying the scenarios beyond 2100 as decisions must be made with awareness of the assumptions about the future being made. These include the level of warming that is assumed and the degree to which low confidence processes are to occur and additionally drive high amounts of future sea level rise.

These ranges in 2150 are among the most likely values to be revised in future updates as scientific understanding and ice-sheet

modeling continues to evolve. In sum, deep uncertainties and ambiguity embedded in the High Scenario frame a worst case as we currently understand it and should be used with caution and consideration of the underlying assumptions in planning adaptation.

2.4. Sea Level Scenario Storylines

2.4.1 Scenario exceedance probabilities

Given that sea level rise is highly dependent on if and how fast the world's nations reduce global emissions and mitigate warming trends, there are no probabilities that can be assigned directly to each of the Sea Level Scenarios. Instead each scenario integrates information on a potential future pathway for warming levels and emissions. By extension, assumptions about future warming levels can be translated into the probability of exceeding a particular Sea Level Scenario in that assumed future (Table 2.2).

The IPCC AR6 Working Group III (IPCC, WGIII, 2022) assessed that, extrapolating current policies for greenhouse gas emissions warming in 2100, global surface temperature warming is projected to be roughly 3°C above pre-industrial levels. This contextual information can be used to construct “storylines” for each scenario, describing what the future will look like in each case. The storylines for Low, Intermediate Low, Intermediate, Intermediate High and High Scenarios are described in more detail below. Notably, the percentages in Table 2.2 shown as <0.1% can be interpreted as effectively zero, although they are subject to specifications in the inputs and framework used to generate the projections.

35. Kopp et al., 2023.

TABLE 2.2. Exceedance probabilities for the Sea Level Scenarios based on IPCC warming level-based GMSL projections. Global mean surface air temperature anomalies are projected for years 2081-2100 relative to the 1850-1900 climatology. Global surface temperatures are currently on track to reach 3.0°C above pre-industrial levels by 2100, assuming current rates of emissions-driven warming. Therefore, any temperature anomalies less than (e.g., 1.5 or 2.0°C) or greater than (e.g., 4.0 or 5.0°C) the current trajectory implies lower or greater rates of warming by 2100, respectively. Low warming in the sixth column broadly refers to temperature anomalies less than 2.0°C, and high warming refers to temperature anomalies greater than 4.0°C. As an example of how this table can be read, the third row could be used to produce the following two sentences: “Assuming 3°C of warming in 2100, there is a 5% chance of exceeding the Intermediate Scenario in 2100” and “Assuming high levels of warming in 2100 and contributions from the low confidence processes, there is a 49% chance of exceeding the Intermediate Scenario in 2100.”

| Global Mean Surface Air Temperature 2081-2100 | 1.5°C | 2.0°C | 3.0°C | 4.0°C | 5.0°C | Low Confidence Processes, Low Warming | Low Confidence Processes, High Warming |
|---|-------|-------|-------|-------|--------|---------------------------------------|--|
| Low Scenario | 92% | 98% | 99.5% | 99.9% | >99.9% | 90% | 99.5% |
| Intermediate-Low Scenario | 37% | 50% | 82% | 97% | 99.5% | 49% | 96% |
| Intermediate Scenario | 0.5% | 2% | 5% | 10% | 23% | 7% | 49% |
| Intermediate-High Scenario | 0.1% | 0.1% | 0.1% | 1% | 2% | 1% | 20% |
| High Scenario | <0.1% | <0.1% | <0.1% | <0.1% | 0.1% | <0.1% | 8% |

2.4.2. Low Scenario

The target of 1 foot of increase in global sea level rise by 2100 is set under the assumption of the current rate of sea level rise continuing on into the future. This assumption is inconsistent with current observations of an acceleration in sea level rise, but could still be considered plausible under the most aggressive emission reduction scenarios. As a result, the Low Scenario provides the lower bound for plausible sea level rise in 2100 and sits below the median value for all AR6 scenarios at all

times between 2020 to 2150. The likelihood of exceeding this Sea Level Scenario is greater than 90% at all warming levels.

SUMMARY:

Aggressive emissions reductions leading to very low future emissions; the scenario is on the lower bounding edge of plausibility given current warming and sea level trajectories, and current societal and policy momentum.

2.4.3. Intermediate-Low Scenario

This scenario arises under a range of both future warming levels and possible SSPs, spanning low, intermediate and high emissions pathways, and integrates many of the AR6 SSP pathways as a result (see Figure 2.2) This scenario is consistent with the median projected sea level rise in a 2°C world, which means there is a 50% probability of exceeding this scenario with 2°C of additional warming by 2100. At a warming level of 3°C in 2100, the probability of exceeding this scenario is 82%. Given the extrapolation of GMSL to 2100 (approximately 2.2 feet³⁶), the current projection of future warming of 3°C, and the range of sea level rise across the IPCC AR6 scenarios (Figure 2.4), the Intermediate Low Scenario provides a reasonable lower bound for the most likely range of sea level rise by 2100. Since the low confidence processes are not important to this scenario, the range of possible sea level rise after 2100 does not expand significantly.

SUMMARY:

A range of future emissions pathways; a reasonable estimate of the lower bound of most likely sea level rise in 2100 based on support from sea level observations and current estimates of future warming.

2.4.4. Intermediate Scenario

The Intermediate Scenario is driven dominantly by high emissions scenarios, and thus higher warming levels. For the first time in the scenarios, the low confidence projections from the IPCC AR6 contribute significantly and provide about 25% of the pathways for reaching the Intermediate Scenario target by 2100. Given the extrapolation of GMSL to 2100 and the range of sea level rise across the IPCC AR6 scenarios (Figure 2.4), the Intermediate Scenario provides a reasonable upper bound for the most likely range of sea level rise by 2100. At a warming level of 3°C in 2100, the probability of exceeding this scenario is 5%. In a very-high emissions future with low confidence processes, there is about a 50% chance of exceeding the Intermediate scenario in 2100.

SUMMARY:

A range of future emissions pathways; could include contribution from low confidence processes. Based on sea level observations and current estimates of future warming, a reasonable estimate of the upper bound of most likely sea level rise in 2100.

36. Nerem et al., 2022.

2.4.5. Intermediate-High Scenario

Pathways combining both higher emissions and low confidence processes become the majority, with over 50% of the samples used to construct this scenario coming from the SSP5-8.5 scenario. At all times from 2020 to 2150, the Intermediate High Scenario exceeds the median value of the AR6 scenarios. This scenario is similar to the high-end estimate from van de Wal et al. (2022) under the assumption of high levels of warming in 2100. At a warming level of 3°C in 2100, the probability of exceeding this scenario is 0.1% when not considering the low confidence processes, emphasizing the degree to which these processes are needed to get to this scenario. With the low confidence processes, the probability of exceeding this scenario is approximately 20% for very high warming levels.

SUMMARY:

Intermediate-to-high future emissions and high warming; this scenario is heavily reflective of a world where rapid ice sheet loss processes are contributing to sea level rise.

2.4.6. High Scenario

Pathways combining both high emissions and low confidence processes are dominant, providing over 80% of the samples to construct the scenario. Low emissions pathways are not plausible under this scenario, and intermediate emissions pathways require a significant contribution from rapid ice sheet loss processes. Before 2100, the High Scenario is significantly above the range of SSP AR6 scenarios, although the range of plausible sea level expands beyond 2150. The probability of exceeding the High Scenario in 2100 is less than 0.1% for all warming levels without considering low confidence processes. With very high emissions and warming and contributions from the low confidence processes, this probability increases to 8%.

SUMMARY:

High future emissions and high warming with large potential contributions from rapid ice-sheet loss processes; given the reliance on sea level contributions for processes in which there is currently low confidence in their understanding, a statement on the likelihood of reaching this scenario is not possible.

3. Combined Impacts of Sea Level Rise and Other Coastal Hazards

LEAD AUTHORS: Dr. Patrick Barnard, Dr. Gary Griggs, Dr. Kristina Hill

SUPPORTING AUTHORS: Dr. Susheel Adusumilli, Dr. Dan Cayan, Laura Engeman, Dr. Benjamin Hamlington, Dr. Felix Landerer, Dr. Phil Thompson, Ocean Science Trust

BOX 9:

Key Takeaways:

Combined Impacts of Sea Level Rise and Other Coastal Hazards

- **Rising seas** will exacerbate coastal hazards such as flooding, coastal erosion, and shallow groundwater, which will impact public infrastructure, private development, livelihoods, public health and safety, and natural systems. In the coming decades, these hazards and their associated social and economic impacts will be the most evident manifestation of future climate change in coastal communities.
- **Over time**, sea level rise will increase the frequency of coastal flooding events, which occur when sea level rise amplifies short-term elevated water levels associated with higher tides, large storms, El Niño events, or when large waves coincide with high tides. California coastal communities need to be aware of and prepared for already increasing high-tide flooding frequency to further accelerate in the early 2030s, as well as continued increase in flood occurrence.
- **Sea level rise** will increase rates of retreat of coastal cliffs and bluffs, the erosion and/or landward migration of beaches, and the loss of coastal wetlands, tidal marshes and sand dunes where barriers exist to their landward migration. Without intervention, approximately 24 to 75% of California's sandy beaches could be seasonally or permanently lost due to sea level rise.
- **In coastal areas** with shallow unconfined groundwater, the water table will rise as sea level rises. This groundwater rise can mobilize soil contaminants, compromise below-ground infrastructure, and may result in saltwater intrusion into freshwater aquifers. The societal impacts of shallow and emerging groundwater are projected to be comparable to overland flooding impacts during the 21st century.
- **Storm events** will become more damaging and dangerous as climate change and sea level rise continue. Impacts from high-intensity coastal storms in combination with sea level rise and high tides include accelerated cliff and bluff erosion, coastal flooding and beach loss, and subsurface contaminant mobilization. Sea level rise will increase the exposure of communities, assets, services and culturally important areas to significant impacts from coastal storms.

IN THE NEAR TERM, the most obvious demonstrations of sea level rise are likely to be increased coastal flooding, shoreline and cliff retreat, groundwater rise, and habitat migration or loss. These coastal processes will be exacerbated by rising seas and will threaten and damage infrastructure and development, impact livelihoods and public health and safety, and jeopardize natural systems.

This chapter provides an overview of coastal hazards that will increase in frequency and/or intensity with rising sea levels, as outlined in Chapter 2.0. The information in this chapter and resources included in Appendix 4 are intended to inform steps 3 and 4 of the stepwise planning process described in Chapter 4.0. Steps 3 and 4 recommend selection and analysis of storm conditions to understand vulnerability to coastal and inland flooding under Sea Level Scenarios for all projects. The stepwise process also recommends consideration of extreme water levels that lead to flooding, coastal erosion, and groundwater hazards on a case-by-case basis. This chapter describes how rising sea levels are expected to affect these coastal processes so that vulnerability assessments can be conducted with an awareness of relevant coastal hazards.

OPC, working in partnership with other state and federal agencies, is committed to continuing research, investments and actions that support further guidance to evaluate and address impacts from sea level rise and other coastal hazards.

3.1. Flooding and Sea Level Rise

Low-lying shoreline areas, which are often heavily populated, are the most affected by coastal flooding at present during extreme high tides and storms, and will be increasingly impacted as sea level rise continues to accelerate. As seen in January 2023 across many locations in California, intense storms can cause extreme coastal water levels, which result in coastal and inland flooding. Increasingly frequent flooding events routinely affect transportation corridors and impact coastal recreational facilities and other infrastructure and cause recurring damage to private and public structures. To date, frequent locations of coastal flooding include the shorelines of Humboldt and San Francisco Bays, Capitola and Rio del Mar on the Central Coast, and Long Beach, Seal Beach, Sunset Beach, Huntington Beach, Newport Beach, and Imperial Beach in southern California.



High water levels along the coast can cause coastal flooding and severe erosion events and are likely to become more damaging as sea level rises. The continuing rise in sea level across California is predicted to lead to an exponential increase in the frequency of coastal flooding events, doubling with approximately every 2-4 inches of sea level rise.³⁷ This translates to flooding that will also last longer, extend further inland, and to greater depths. Today's once-in-a-lifetime coastal flood could occur annually by 2050 and daily by 2100.³⁸ Disruption of highway and rail traffic, and other transportation corridors will have cascading impacts across our communities, and will be among the most significant impacts affecting the largest number of people and threatening public safety.

High-tide flooding occurs when high tides, often coupled with other factors that raise the local or regional water level, including wave runup, storm surge, seasonal variations, and regional sea level, cause flooding. The frequency and severity of high-tide flooding is projected to increase as sea levels rise. At present, we can get a glimpse into the future during king tides – extreme high tides that happen several times a year – as higher water levels will occur more frequently in the future as sea level continues to rise.

High-tide flooding can vary across locations – even from one community to the next – depending primarily on coastal topography and the elevation of public infrastructure and private development. A statistical model developed by Thompson et al. (2021) projects how the frequency of high-tide flooding might increase during the 21st century at fourteen NOAA tide gauges in California under the different Sea Level Scenarios (see Box 10) below for more information on this tool). Using the definition of minor high-tide flooding established by NOAA,³⁹ the frequency of high-tide flooding in California is expected to increase significantly during and after the 2030s, beyond the increase in frequency already evident. Under the Intermediate Scenario, the frequency of minor high-tide flooding is projected to increase by a factor of three to four depending on location from 2030 to 2050.

In addition, California coastal communities will also be exposed to more frequent high-tide flooding events due to occurrences of the El Niño phase of the El Niño Southern Oscillation and other oceanic events. Southern California shorelines have a greater than 30% chance of experiencing at least 20 NOAA Minor high-tide flooding days during a single year by 2030, with San Diego having the highest probability at almost 60%.⁴⁰

37. Vitousek et al., 2017.

38. Taherkhani et al., 2020.

39. <https://oceanservice.noaa.gov/facts/high-tide-flooding.html>

40. Thompson et al., 2021.



BOX 10:

High-Tide Flooding

The NASA Sea Level Change Team, in partnership with the University of Hawai'i Sea Level Center, developed a [Flooding Analysis Tool](#) that provides observed and projected analysis of high-tide flooding days for 14 tide gauge locations in California to help understand and determine the impacts of future high-tide flooding. The high-tide flooding projections in the Flooding Analysis Tool incorporate the same sea level scenarios that were downscaled for this report, so the tool can be used to determine how high-tide flooding has increased and how it may increase by location during the 21st century under the different Sea Level Scenarios.

The information provided by the Flooding Analysis Tool is site-specific for each tide gauge but is not extrapolated for areas beyond the tide gauges. If a tide gauge does not exist at the desired location, analysis from the closest tide gauge can provide useful information, but it is important to consider the potential impact of local factors that can differ over short distances that may impact high-tide flooding. The information provided

by the Flooding Analysis Tool may be helpful in determining local and regional vulnerability to high-tide flooding into the future, informed by past experiences with high-tide flooding events and projected sea level rise. While evaluation of projected high-tide flooding days is not included as part of the stepwise process in Chapter 4.0, it may be important to include in vulnerability assessments on a case-by-case basis, particularly for locations that have previously been impacted by high-tide flooding.

The Flooding Analysis Tool can be used to view high-tide flooding projections. The About tab provides instructions for how to use the tool, including selection of location, flood threshold, and sea level scenario. The Observed Flooding tab allows users to determine relevant flood thresholds based on the known impacts of past high sea level events. Based on the selected combination of location, threshold, and scenario, the Projected Flooding tab will show the expected number of flooding days per year during the 21st century.

BOX 10 CONT.




FIGURE 3.3. Tables produced through the Flooding Analysis Tool showing how the frequency of flooding will evolve in the near term at four California tide gauges. The La Jolla, Los Angeles, and Humboldt Bay tables are based on NOAA’s Minor flooding threshold under the Intermediate Sea Level Scenario, while the San Francisco table uses NOAA’s 2-year flood threshold under the Intermediate Sea Level Scenario because that threshold corresponds to recent flooding events. Projected high-tide flooding days and analyses for the other California tide gauges under a range of NOAA flooding thresholds can be found through the Flooding Analysis Tool.

41. <https://sealevel.nasa.gov/flooding-analysis-tool/instructions?station-id=9414290>

By 2050, almost every California tide gauge location has a greater than 50% chance – and most have a greater than 75% chance – of experiencing at least one year with 30 minor high-tide flooding days and at least a single month with 10 minor high-tide flooding days. Impacts associated with these minor but increasingly frequent flooding events, such as a brief disruption to transportation systems and economic activity, will have cumulative impacts that can be greater than infrequent extreme events.⁴²

California coastal communities need to be aware of and prepared for already increasing high-tide flooding frequency to further accelerate in the early 2030s. Statewide population exposure to daily coastal flooding varies by a factor of 12 from the Low Sea Level Scenario around 2050 (i.e., 0.25 m SLR = 38,000 residents, expected around 2050 for other SLR scenarios) to the High Sea Level Scenario around 2100 (i.e., 2.0 m SLR = 440,000 residents, high scenario at 2100) for everyday (no-storm) conditions.⁴³

Severe storms often bring heavy precipitation resulting in higher stream flows, and strong winds and low atmospheric pressure that generate large waves and storm surge, which can have a compounding effect on shoreline flooding. Compound flooding events often have greater impacts than isolated hazards, like high tides and river flooding that occur asynchronously. As sea level rises incrementally, more frequent and severe compound flooding events may be one of the most obvious and damaging effects of sea level rise.⁴⁴

BOX 11:

New research underway to understand coastal flooding and extreme water levels

As already experienced in California, heavy rainfall events can coincide with ocean-driven coastal waves and storm surge leading to a compound flooding hazard. These co-occurrences of wave overtopping and heavy runoff often lead to significantly more damages from flooding and shoreline erosion. With research funding from the California Energy Commission (CEC), researchers from Scripps Institute of Oceanography are advancing our understanding of, and ability to project, extreme high water events.

Using historic sea level elevations and the maximum sea level heights (99.9 percentile) seen at specific tide gauges in California, often during El Niño seasons, this research will provide hourly projections of how often these maximum sea level records are expected to be reached and exceeded in the future. These projections are consistent with the scenarios included in this report, but provide a different framing to emphasize the expected increase in frequency of today's extreme sea level events as we reach higher levels of sea level rise.

In addition, Scripps researchers are developing a combined model projecting the likelihood of extreme sea level events co-occurring with extreme precipitation storms. Researchers will produce high-resolution climate models to identify unusually high precipitation that additionally contributes to coastal flooding. This research is expected to help inform the state's understanding of how storm events are expected to drive significant coastal runoff and coastal wave overtopping leading to greater flooding frequency, extent, depth, and magnitude of damages.

42. Mofstakhari et al., 2017; Ghanbari et al., 2021.

43. Barnard et al., 2019.

44. Singh et al., 2023.

3.2. Groundwater and Sea Level Rise

3.2.1. Saltwater Intrusion

Increased demand for groundwater in many of California's fertile coastal plain areas, both for agricultural and domestic uses, has led to excessive pumping, which has lowered the groundwater table, and allowed intrusion of seawater into aquifers, referred to as saltwater intrusion. Because saltwater has a greater density than freshwater, it moves into freshwater aquifers in the form of a wedge, with greater intrusion at depth than at the surface. Saltwater intrusion is a concern in coastal California even before factoring in sea level rise, but sea level rise can expedite the landward moment of saltwater wedges by forcing the saltwater wedge further inland.

Saltwater intrusion is a problem because when salt or brackish water appears in wells, it can render them unusable for drinking and agriculture.⁴⁵ Increases in groundwater salinity and higher water tables can also impact building foundations, infiltrate sanitary sewers and stormwater pipes, and corrode underground utilities that were not designed to be submerged in saltwater.⁴⁶ This puts foundations, underground sewer pipes, and public utility infrastructure (e.g., gas pipelines, fiber optic cables, electrical lines, sewage pump stations) at progressive risk of failure. Saltwater intrusion in the lower Salinas Valley around southern Monterey Bay has been documented since 1944 and now extends eight miles inland nearly to the city of Salinas (Figure 3.4). Saltwater intrusion has also been a major problem in the Sacramento Delta area, in Santa Clara County adjacent to San Francisco Bay, and on the Oxnard Plain in Ventura County. Adaptation planning should consider the risk that saltwater intrusion into groundwater could increase significantly if pumping occurs over this saltwater wedge.⁴⁷ Drawing saltier groundwater closer to the surface can cause corrosion impacts and shorten the useful life of pumps.

45. Jakovic et al., 2016.; Loáiciga et al., 2012.

46. Habel et al., 2023.

47. Jakovic et al., 2016.




FIGURE 3.4. Historical progression of seawater intrusion into the aquifer in the lower Salinas Valley used for domestic and agricultural uses. Map Source: Monterey County Water Resources Agency.⁴⁸

3.2.2. Groundwater rise

Along shorelines, subsurface saline water will penetrate further inland as the ocean rises, pushing less-dense freshwater upward. This will raise the water table and increase the potential for coastal groundwater hazards. Low-lying coastal areas of California with shallow unconfined aquifers occur along the shoreline, such as the margins of Humboldt Bay, San Francisco Bay, Monterey Bay and Santa Monica Bay, and San Diego Bay, are particularly vulnerable to these groundwater hazards. Rising groundwater can reduce domestic and agricultural water quality, detrimentally impact underground infrastructure (e.g., foundations,

basements, and utility lines), and flood low-lying areas along the coast or tidally influenced waterways. The response of the water table is dependent on the distance from the coast, topographic features, the magnitude of sea level rise, the permeability of the subsurface materials, and rainfall.

Recent studies show that coastal communities will be much more exposed to shallow coastal groundwater hazards over the next century, defined here as the water table within 6.5 feet (2 m) of the ground surface, than overland flooding alone. Hazards from shallow groundwater have the potential to affect 1.9 million California residents under

48. Monterey County Water Resources Agency, 2013.

the Intermediate Sea Level Scenario (i.e., 3.3 feet or 1 m), nine times higher than overland coastal flooding for the same scenario.⁴⁹ Preliminary research indicates potential serious risks for human and ecosystem health near urban sites that handle hazardous materials and sites that contain soil contaminants.⁵⁰ Subsurface contaminants in the thousands of buried waste disposal sites surrounding San Francisco Bay will be mobilized by rising sea level and groundwater. Mobilization of soil toxins can occur as a result of a gradual increase in groundwater elevation from sea level rise or can happen during a severe storm event that causes flooding and rapid groundwater rise.

A recent study found that groundwater rise may affect thousands of contaminated sites in the San Francisco Bay Area, and that socially vulnerable communities are disproportionately exposed to this risk.⁵¹ Although the effects of sea level rise on contaminated sites across the rest of coastal California have not been thoroughly studied, contaminant mobilization is a concern anywhere such sites exist near the coast. Over time, with a continuing rise in sea level, shallow groundwater rise will gradually shift further inland, exposing new sites that were designed without consideration for shallower, more saline, or emerging water tables.

An approximately linear relationship can exist between sea level rise and groundwater rise within as much as half a mile of a shoreline or bayline if there is limited surface drainage. In other words, the maximum possible projected future groundwater rise can be estimated by adding the amount of projected sea level from the Sea Level Scenarios to current groundwater levels.⁵² This approximate one-to-one relationship between groundwater rise and sea level rise can be used in areas within approximately a half-mile from the shoreline that do not have precise regional measurements or projections. Farther from the shoreline, groundwater rise tapers and will not rise in a 1:1 relationship to sea level. In areas where groundwater pumping occurs or where rising groundwater intersects surface drainage features, the response of the water table to sea level rise will be damped or reduced.

In some areas, the water table may rise enough to emerge at the surface, which can cause temporary flooding and/or permanent inundation. Even if water tables do not become emergent, groundwater can still be problematic before reaching the surface by impacting underground infrastructure. It can infiltrate storm drains, causing spot flooding; destabilize pipes that were not designed for frequent subsurface water exposure; undermine foundations; and corrode infrastructure not designed for exposure to saline groundwater.

49. Befus et al., 2020.

50. Cushing et al., 2023; Hill et al., 2023.

51. Cushing et al., 2023; Hill et al., 2023.

52. Befus et al., 2020.

53. Habel et al., 2023.

BOX 12:**Screening for exposure to groundwater rise**

Our Coast, Our Future (OCOF), developed by Point Blue Conservation Science and USGS Pacific Coastal and Marine Science Center, is a user-friendly web-based tool for understanding and viewing potential flooding, coastal erosion, and shallow groundwater hazards based on model outputs from the Coastal Storm Modeling System (CoSMoS). This tool allows users to identify areas along California's coastline that will likely be exposed to, and thus may be at risk from, these coastal hazards due to projected sea level rise.

As it specifically relates to groundwater rise, this tool can be used to assess where coastal assets and resources may be exposed to shallow groundwater. By incorporating the sea level scenarios for a given location from this guidance (see Appendix 2), users of this tool can conduct a rapid assessment of

where communities and infrastructure may be susceptible to shallow groundwater hazards.

Additionally, OCOF can also be used to identify areas of groundwater rise (i.e. shoaling) and flooding (i.e. groundwater emergence). By incorporating an area's respective Sea Level Scenarios, OCOF depicts areas of varying groundwater hazards, including: (1) marine inundation (i.e. coastal ground surface below MHHW sea level), (2) emergent (i.e. water table reaches ground surface), (3) very shallow (i.e. water table within 3.3 ft of ground surface), (4) shallow (i.e. water table within 3.3 - 6.6 ft of ground surface), and (5) moderate (i.e. water table within 6.6 - 16.4 ft of ground surface). These levels represent varying risks of shallow, subsurface infrastructure (e.g., basements, foundations, gas and water lines, septic systems, etc.) from groundwater hazards. An example for Richmond, CA is provided below (Figure 4.3). With this information in hand, practitioners can better assess the need to collate, or invest in collecting, additional data and information to quantify risk and or potential impacts.




FIGURE 3.5. Snapshot output from OCOF illustrating groundwater hazards for Richmond, CA using the Intermediate Scenario for 2050 (0.8 ft).

3.3. Coastal and Shoreline Erosion

3.3.1. Loss or Migration of Beaches

Beaches serve as the first line of defense for many coastal communities against storm-driven flooding because they dissipate wave energy. Beaches with no back beach obstructions will tend to migrate landward as sea level rises and/or sediment supply is reduced. However, if landward migration is inhibited by physical barriers, such as resistant cliffs, seawalls, rock revetments, railway lines, highways, or other urban development, beaches will narrow or possibly drown, being permanently inundated between a rising ocean and the resistant feature, a phenomenon known as coastal squeeze or passive erosion.

The response of California's sandy shoreline to sea level rise and storms has been projected across the entire state using the CoSMoS-COAST shoreline model.⁵⁴ This model integrates local information on beach behavior based on satellite observations from the last 25 years (1995-2020), and then makes projections of shoreline change based on wave-driven longshore and cross-shore transport of sand due to future wave conditions and sea level rise. This model also makes projections considering both the existence and removal of protective shoreline structures such as seawalls or revetments. Across the range of the sea level scenarios, and without preventative or management action, 24 to 75% of sandy beaches could be seasonally or permanently lost by 2100. Changes in sediment supply could reduce or increase this loss; for example, dams have reduced the natural sand supply to California beaches from rivers and streams by 23%, including a 50% reduction in Southern California, since the late 19th century.⁵⁵

3.3.1. Loss or Migration of Beaches

The great majority of California's 1,100-mile coast (790 miles or 72%) consists of actively eroding sea cliffs. About 650 miles of this is composed of lower-relief cliffs and bluffs typically eroded into uplifted marine terraces. Many of the state's communities are built on these features - including Fort Bragg, Mendocino, Pacifica, Half Moon Bay, Santa Cruz, Pismo Beach, Santa Barbara and many of the cities in southern Orange and northern San Diego Counties. Most of these cliffs and bluffs are actively eroding, although 38% of the coast of Southern California has now been armored.⁵⁶ As sea levels continue to rise, waves will reach and impact the base of coastal cliffs, bluffs, and dunes more often which will lead to increased erosion rates and threats to blufftop development and infrastructure as shown in Figure 3.6.

54. Vitousek et al., 2017; 2021; 2023.

55. Slagel and Griggs, 2008.

56. Griggs and Patsch, 2019.



FIGURE 3.6. *Bluff retreat in Pacifica ultimately led to the demolition of these apartments.*
Photo Source: Joel Avila / Hawkeye Photography.

Historical cliff retreat rates in California range from several centimeters to a meter or more per year.⁵⁷ Cliff recession rates vary based on a number of regional and local factors, including wave impacts at the base of the cliff, tidal range, rock type, joint density, tectonic forces, protection structures, drainage and groundwater flow, climate variability, and storm frequency. However, regardless of these variables, an increase in sea level will result in greater exposure of the cliff base to wave impacts, and the rate of cliff retreat is likely to increase.

Limber et al. (2017) projected statewide cliff retreat for the five Sea Level Scenarios. The results show an increase in the rate of retreat over historical rates with continued sea level rise, and as much as a doubling with two meters of sea level rise. For example, under the Intermediate Scenario of 1 meter by 2100, cliffs are projected to retreat 23 m (75 ft), on average, by 2100, although retreat rates will vary widely depending upon local cliff properties and wave energy. Nonetheless, both protected and unprotected cliff top communities will be at an even greater risk over the 21st century. Each coastal community will need to undertake vulnerability assessments to determine which facilities and development are at greatest risk and then agree on how and when to respond or adapt to expected future conditions.

57. Hapke and Reid, 2007; Swirad and Young et al., 2022; Griggs, et al., 2005.

3.3.3. Loss or Migration of Coastal Ecosystems and Species

Coastal ecosystems will be significantly impacted by sea level rise across the state. Habitats such as beaches, dunes, and tidal marshes will be exposed to more frequent flooding, erosion and coastal squeeze, although in undeveloped areas where there are no barriers, they will continue to migrate inland, as they have historically. In many locations, the habitat space for many species will be progressively constricted between rising seas and urban development, restricting landward migration.⁵⁸ Hard armoring along the coast, including rock revetments and seawalls, contributes to habitat loss because rising sea levels submerge coastal habitats while these artificial structures prevent landward migration.

Coastal wetland elevation has kept pace with lower rates of sea level rise throughout the holocene due to sediment accumulation and available accommodation space.⁵⁹ The ability of tidal wetlands and marshes to continue to build elevation at a sustainable rate relative to higher sea levels of the future depends primarily on the rate of sea level rise, sediment deposition,⁶⁰ building of below ground peat soils by wetland vegetation⁶¹ and decomposition and sediment compaction rates. Under higher sea level scenarios, Thorne et al. (2018) predict that substantial vegetated tidal marshes in California will transition to mud flats by the end of the 21st century due to constraints on horizontal migration in urbanized estuaries (San Francisco Bay, for example). The historical alteration of watersheds and tidal marshes across the state and reductions in freshwater and sediment supply that reduce vertical growth potential will further challenge marsh sustainability and the associated ecosystems under sea level rise. Loss of coastal wetlands and tidal marshes would also mean losing the ecosystem services they provide, including maintaining water quality, sequestering carbon, reducing turbidity, and providing buffers against storm waves and flood waters.⁶²

Species that depend upon coastal habitats are also at risk from sea level rise. Increased inundation and higher salinity will alter shoreline plant distributions, which will have outside impacts on coastal fauna.⁶³ A 2018 study found that eight imperiled species only occur in areas that are projected to be inundated with five feet of sea level rise, including coastal dunes milk-vetch, California seablite, and California Ridgway's rail.⁶⁴ Documented haul-outs and breeding colonies for Pacific harbor seals and Northern elephant seals, which are critical to maintaining populations, are

58. Dugan et al., 2008; Dugan and Hubbard, 2010; Myers et al., 2019; Barnard et al., 2021.

59. Kirwan and Megonigal, 2013.

60. Leonard, 1997; Buffington et al., 2021.

61. Nyman et al., 2006; Cherry et al., 2008.

62. Barbier et al., 2013.

63. Goals Project, 2015.

64. Heady et al., 2018.

also highly vulnerable to rising seas. Nesting habitats for shorebirds like California least tern, black oystercatcher, and Western snowy plover may be inundated without opportunity for landward migration in locations where beaches are bordered by urban environments, cliffs, or bluffs. Black oystercatchers and Pacific harbor seals rest on beach and rocky intertidal habitats, seeking areas that are free from predators and human disturbance. The small pocket beaches and rocky intertidal areas at the base of cliffs that these species typically select are particularly vulnerable to sea level rise.

3.3.4. Threats to Coastal Access and Recreation

As beaches are lost to rising seas as described above, space available to Californians and visitors for recreation along the coast decreases. Losing beaches to sea level rise in itself constricts available space for recreation, but beach accessibility is also contingent on the infrastructure and amenities that support visitation and recreation.⁶⁵ Coastal access features like trails and stairways, parking facilities, lifeguard towers, and amenities like restrooms, showers and picnic areas all contribute to the accessibility of coastal areas. As sea levels rise, these features will become vulnerable to inundation and damage from coastal storms. Beachgoers have different preferences for features, facilities, and amenities at coastal access sites,⁶⁶ so loss of infrastructure will not affect all beachgoers equally. For example, loss of restroom amenities will likely be more impactful on visitors who must travel further to reach the beach and loss of lifeguard towers will likely deter individuals with less confidence as swimmers; underlying each of these cases are significant issues relating to equity and environmental justice as beach access becomes increasingly challenging. An online geodatabase provides a statewide picture of how coastal access locations overlap with rising seas.⁶⁷

Rising seas will also have impacts on surf breaks, which hold substantial recreational value in California. As sea levels rise, water depth at current surf break locations will increase. Deeper water means either that only larger waves can break at that precise location, or that smaller waves will break closer to the present shoreline in shallower water.⁶⁸ As shoreline geography and development allows, some surf breaks will migrate landward but most are likely to meet a suite of challenges, imperiling a valued recreational resource.⁶⁹

65. Patsch and Reineman, 2023 (accepted).

66. Christensen and King, 2017.

67. <https://www.arcgis.com/apps/dashboards/19ac80fe57e747ac9caaf966b29cb9c4>. Note: While this dataset can provide valuable insights into potential vulnerabilities, it should not be used as the sole basis for critical decisions related to coastal management, emergency response, or infrastructure planning.

68. Reineman et al., 2017.

69. Sadrpour and Reineman, 2023.

3.4. Preparing for Extreme Coastal Storms

Although the near-term and cumulative effects of sea level rise described above occur gradually over time, when extreme coastal storms occur coincident with higher background sea levels, very high or king tides and/or short-term sea level rise (such as during an El Niño event), the impacts and damages can be sudden and severe. For example, extreme coastal storms with large waves can cause widespread flooding in low-lying areas, extensive beach loss, and cliff or bluff collapse (Figure 3.4). As a first pass at understanding coastal risk, communities need to apply the history of damage from past winter storm events (January 2023, for example) including during powerful El Niños (1982-83, 1997-98, and 2015-16) to plan in advance for how to respond and adapt to similar events in the future in order to limit losses and disruption, all of which will be exacerbated with a higher baseline of sea level. Another approach is to refer to the results of modeled extreme storm scenarios, such as the annual, 20-year, and

100-year storm events, coupled with sea level rise, to more broadly understand the risks of climate-driven coastal hazards.

CoSMoS⁷⁰ can be used to understand the cumulative impacts of sea level rise and storms on California coastal communities. Adding storms increases population exposure by 50-340% over daily high tide conditions across the range of Sea Level Scenarios presented in Chapter 2.0. Approximately 700,000 California residents and \$250 billion in property could be exposed to flooding by 2100 under the High Scenario and a 100-year storm.⁷¹ The San Francisco Bay Area accounts for two-thirds of future flood risk of the California population and property values state-wide, but low-lying areas across the state are also at risk, including numerous coastal and estuarine communities, as well as airports, port facilities, transportation corridors, and public utilities

70. <http://www.usgs.gov/cosmos>

71. Barnard et al., 2019.

FIGURE 3.7. *Damage to West Cliff Drive in Santa Cruz during high tides and extreme waves in January 2023.*





4. California Sea Level Rise Policy Guidance

AUTHOR: Ocean Protection Council

4.1. Summary

IN THIS UPDATE, which replaces the 2018 California Sea-Level Rise Guidance, California continues its commitment to science-based policy and decision-making. This revised policy guidance incorporates the science update presented in Chapters 2.0 and 3.0 to ensure that state, regional, and local sea level rise adaptation planning and project decisions are consistent and grounded in the best available science. As in 2018, this guidance is deliberately structured to be both precautionary and flexible to accommodate local and regional priorities and the broad array of decision-making contexts in which planning for sea level rise is relevant.

In Chapter 2.0, five Sea Level Scenarios are constructed and presented for the California coast. Table 4.1 shows the statewide average values for these five scenarios which use a single average value of vertical land motion corresponding to a negligible rate of 0.10 mm/year uplift. Sea Level Scenarios are also presented for the 14 tide gauges in Appendix 2

and deviations from the statewide average are due to differences in localized vertical land motion (uplift or subsidence).

The Sea Level Scenarios show greater certainty in the amount of sea level rise expected in the next 30 years than previous reports and demonstrate a narrow range across all possible emissions scenarios. Statewide, sea levels are most likely to rise 0.8 ft (Intermediate Scenario) by 2050. In the mid-term (2050-2100), the range of possible sea level rise expands due to more uncertainty in projected future warming from different emissions pathways and certain physical processes. By 2100, sea levels are expected to rise between 1.6 ft (Intermediate-Low) and 3.1 ft (Intermediate Scenario), although higher amounts cannot be ruled out. Beyond 2100, the range of possible sea level rise becomes increasingly large due to uncertainties associated with physical processes, such as earlier-than-expected ice sheet loss and resulting future sea level rise. Statewide, sea levels may rise from 2.6 ft to 11.9 ft (Intermediate-Low to High Scenarios) by 2150, and even higher amounts are possible.

TABLE 4.1. Median values for Sea Level Scenarios for California, in feet, relative to a 2000 baseline. These statewide values all incorporate an average value of vertical land motion corresponding to a negligible rate of 0.1 mm (0.0003 ft) per year uplift. Evaluation of the Intermediate, Intermediate-High and High Scenarios (outlined in red below) is recommended to inform appropriate sea level rise planning and project decisions.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 |
| 2030 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 |
| 2040 | 0.4 | 0.5 | 0.6 | 0.7 | 0.8 |
| 2050 | 0.5 | 0.6 | 0.8 | 1.0 | 1.2 |
| 2060 | 0.6 | 0.8 | 1.1 | 1.5 | 2.0 |
| 2070 | 0.7 | 1.0 | 1.4 | 2.2 | 3.0 |
| 2080 | 0.8 | 1.2 | 1.8 | 3.0 | 4.1 |
| 2090 | 0.9 | 1.4 | 2.4 | 3.9 | 5.4 |
| 2100 | 1.0 | 1.6 | 3.1 | 4.9 | 6.6 |
| 2110 | 1.1 | 1.8 | 3.8 | 5.7 | 8.0 |
| 2120 | 1.1 | 2.0 | 4.5 | 6.4 | 9.1 |
| 2130 | 1.2 | 2.2 | 5.0 | 7.1 | 10.0 |
| 2140 | 1.3 | 2.4 | 5.6 | 7.7 | 11.0 |
| 2150 | 1.3 | 2.6 | 6.1 | 8.3 | 11.9 |

4.2. Stepwise Process to Apply Sea Level Scenarios in Planning and Projects

The following steps provide a decision framework that can be used to guide selection of appropriate Sea Level Scenarios for specific planning and project application. **For the purposes of this guidance, planning refers to local and regional land-use and community planning efforts and projects refer to site-specific siting, conceptual and engineering design, permitting, or construction related to coastal habitats, development, access, or infrastructure. A vulnerability assessment (or similar hazard analysis) is considered to be a step in a planning effort or project rather than a standalone effort.**

BOX 13:**Existing and Ongoing Efforts**

Existing and ongoing efforts to analyze and plan for the impacts of sea level rise should be able to proceed with little to no adjustments; there are likely very few, if any, situations in which work will need to be redone as a result of the new Sea Level Scenarios. Instead, a qualitative evaluation should be performed that compares the sea level rise values used in previous and/or ongoing efforts with the new values and recommendations in this updated Guidance. For instance, existing vulnerability assessments, plans, and projects should be evaluated to determine whether appropriate sea level rise was included (as compared with the new Sea Level Scenarios). Values will likely not match exactly, but the range of new Sea Level Scenarios should be captured. Anticipated impacts will likely remain the same when using existing vulnerability assessments; however, the time horizon at which impacts are expected to occur may shift farther into the future. Adaptation planning using existing vulnerability assessments should adequately reflect that shift.

It is important to note that existing community-level vulnerability assessments may not be sufficient to characterize the site-specific vulnerability necessary to support evaluation of project feasibility or design; in such cases, a site-specific vulnerability assessment should be conducted using the new Sea Level Scenarios included in this guidance.

All vulnerability assessments and adaptation plans should be considered living documents that may merit updates based on evolving scientific understanding of the drivers and pace of sea level rise. To avoid a constant cycle of analysis and planning, the Adaptation Pathways approach (see Box 14) is recommended.

>> STEP 1: *Identify the nearest tide gauge*

>> STEP 2: *Evaluate planning and/or project time horizon(s)*

>> STEP 3: *Choose multiple Sea Level Scenarios for vulnerability assessment*

>> STEP 4: *Conduct vulnerability assessment*

>> STEP 5: *Explore adaptation options and feasibility*

>> STEP 6: *Select phased adaptation approach and/or implement project*

FIGURE 4.1. *The following steps, outlined in the figure and in more detail below, provide a decision framework to guide selection of appropriate Sea Level Scenarios for specific planning and project application.*

>> STEP 1: *Identify the nearest tide gauge*

In California, differences in relative, local sea level rise values (i.e. at tide gauge locations) are due to differences in vertical land motion resulting from local factors such as tectonic uplift and subsidence. A location experiencing tectonic uplift will experience lower observed rates of sea level rise than a location experiencing subsidence.

Appendix 2 provides local sea level projections, which incorporate estimates of local vertical land motion, for each of the 14 NOAA coastal tide gauges, including tide gauges in Alameda and Port Chicago that were not included in the 2018 California Sea-Level Rise Guidance, thus providing more locally relevant information for planning in San Francisco Bay. If the project is nearly equidistant between two tide gauges, it can be appropriate to average the two tide gauges. However, it is important to recognize that the tide gauge Sea Level Scenario tables are site-specific information and it is possible that it will not be representative of localized uplift or subsidence. The vast majority of California's coast and shoreline is not experiencing significant uplift or subsidence that would cause an adjustment from a nearby tide gauge, but in some cases, this should be considered when deciding on a tide gauge table or how/if to average. If an area is experiencing little to no vertical land motion, it is appropriate to use the statewide average table.

Alternatively, when technical capacity and available data allows, jurisdictions that have more localized or site-specific data on vertical land motion can choose to combine the statewide average Sea Level Scenarios in Table 4.1 with their local measurements of vertical land motion for any location on the California coast and San Francisco Bay (see Appendix 3). This can be particularly useful for site-specific projects that are experiencing localized tectonic uplift or subsidence.

>> **STEP 2: Evaluate planning and/or project time horizon(s)**

The time horizon refers to how long a given planning effort or project is intended to function and impacts the selection of the sea level rise value greatly. Many of the planning and projects utilizing this guidance will have time horizons to 2100 or beyond, though some projects may be for shorter-term or temporary development. Planning efforts may also consider both long-term goals and the shorter-term actions

necessary to achieve them. For instance, community visioning and planning (such as for a Local Coastal Program) often considers long-term horizons along with the near-term priority actions the community will take. This approach of phased adaptation, or adaptation pathways (see Box 14), for planning and projects can be the most practical and economical approach in many situations, and ensures appropriate actions are taken as sea levels rise over time. In particular, phased adaptation planning should be considered for adaptation of coastal habitats, which are expected to migrate in response to changing conditions. Evaluating the number and timing of adaptation phases within a longer-term timeline will depend on the specifics of the planning or project effort. However, in some situations, it can make sense to plan and adapt for the entire time horizon, such as for critical infrastructure relocation, or when funding and opportunity allow for it.

Alternatively, rather than using time to identify adaptation phases, it is equally valid to choose sea level rise values (step 3) to correspond to adaptation phases so long as those values roughly capture the time horizon in question. For instance, a community planning effort might choose to develop adaptation pathways for 0.8 feet, 3.3 feet and 5.7 feet sea level rise, which roughly correspond to time horizons of 2050, 2100, and 2150, if using the Intermediate Scenario and depending on location. Since phased adaptation, such as adaptation pathways, is often dictated by thresholds and triggers related to sea level rise and not time, this can be a more practical and straight-forward approach. In this situation, it is necessary to identify the time frames at which water levels are likely to be reached and prepare accordingly, including what would need to be done if sea levels rise faster than anticipated. Many climate change adaptation measures require long lead times to accommodate planning, design, permitting, and implementation. Phased adaptation pathways provide a framework for identifying appropriate suites of action at different climate change thresholds and create a mechanism for addressing uncertainty and allowing for flexibility over time.

BOX 14:**Adaptation Pathways**

Adaptation pathways is the recommended approach for sea level rise adaptation planning and is particularly well-suited for mixed-use and community planning efforts. Site-specific implementation projects can also benefit from adaptation pathways when it is not feasible to adapt to higher levels of sea level rise in the near-term. Adaptation pathways allow decision makers to phase short and long-term adaptation strategies over time to build resilience in the face of uncertainty.

Adaptation pathways, which involves sequencing adaptation actions throughout the lifespan of a project, can facilitate cost-effective near-term implementation while planning for future needs that will be triggered by predetermined thresholds or tipping points. Thresholds, or triggers, may be defined by observed sea level rise or other impacts such as flooding extent and frequency or cost to repair/replace damaged built or natural assets. Actions are typically agreed in advance, for example through collaborative community co-design. Figure 4.2 below shows a hypothetical adaptation pathway derived from the Baylands Goals Project 2015. In this example, decisions are triggered at certain thresholds of sea level rise.




FIGURE 4.2. Hypothetical adaptation pathway showing conceptual phasing of measures triggered by sea level rise. Figure source: San Francisco Bay Shoreline Adaptation Atlas.⁷²

Adaptation pathways provide greater flexibility and opportunities to implement no-regrets actions (e.g. habitat restoration or mitigation) to address near-term impacts, while planning for more costly efforts to address and avoid greater damages in the future.

72. SFEI and SPUR, 2019.

>> **STEP 3: Choose multiple Sea Level Scenarios for vulnerability assessment**

A vulnerability assessment is a key step in building coastal resilience and allows decision makers to evaluate the vulnerability of people, natural resources and infrastructure under various future conditions, as well as their level of comfort with over or underestimating sea level rise. Given the uncertainty in the amount of future sea level rise, it is critical to analyze a range of Scenarios to understand what is at risk. **The most precautionary approach, when feasible, is to evaluate Intermediate, Intermediate-High, and High Scenarios to assess a spectrum of potential impacts, consequences, and responses.** However, in practice, evaluation of all three Scenarios might not be necessary and/or may be cost prohibitive. For instance, for near-term actions (between now and 2050), only the Intermediate Scenario is recommended for analysis and selection in Step 6.

The Low Scenario is scientifically plausible but only with accelerated development of carbon capture technologies and global policy and socioeconomic changes that significantly reduce greenhouse gas emissions. The Intermediate-Low Scenario provides a reasonable estimate of the lower bound for the most likely sea level rise by 2100; however, assuming 3°C of warming in 2100, there is an 82% chance of exceeding the Intermediate-Low Scenario in 2100. California is taking significant action to achieve the state's ambitious clean energy goals and is committed to addressing and mitigating the impacts of climate change. However, to ensure precautionary sea level rise planning and projects that protect public health and safety, the environment, critical infrastructure, and public access, for the purposes of this guidance, the Low and Intermediate-Low Scenarios are not recommended for planning or projects.

The High Scenario is considered to be sufficiently precautionary, even for the most risk averse applications; assuming global surface temperatures reach 3.0°C above pre-industrial levels by 2100, there is an effectively zero percent chance of exceeding the High Scenario in 2100. However, it is still considered plausible and assuming high levels of warming (greater than 4°C) in 2100 and contributions from the low confidence processes, there is an 8% chance of exceeding the High Scenario in 2100; therefore, it is still recommended to consider as part of a worst case analysis, which is particularly important for projects and planning applications with extreme risk aversion. For applications with medium to high risk aversion (as described in more detail in Box 18 below), it may be appropriate to make the Intermediate-High scenario be the highest scenario included in technical analyses and vulnerability assessments (Table 2.2).

Consideration of storm conditions in combination with Sea Level Scenarios is also recommended to evaluate extreme water levels, as appropriate. This will provide information on areas that could temporarily flood during episodic storm events before those areas are permanently inundated with higher sea levels. Storm surge and wave processes (i.e., wave set-up and runup) are the primary contributors to extreme coastal water levels during storms. **For most applications, analysis of 100-year storm conditions is**

recommended with wave-driven processes and storm surge being the most important components to consider. However, the coastal water levels resulting from certain combinations of sea level rise and storm scenarios may be similar to others, and therefore not all scenarios need to be thoroughly evaluated. For instance, across California the Intermediate Scenario with 100-year storm conditions, is very similar to the exposure created by the Intermediate-High Scenario with no storms (although they will differ in inundation versus temporary flooding exposure). It is therefore recommended to consider the consequences of storm-induced extreme water levels on a project-by-project basis. Some online visualization tools (such as Our Coast Our Future, see Box 12) consider the primary drivers of extreme water levels and can be used as a screening tool to help determine what combinations of Sea Level Scenarios and storm conditions are appropriate for analysis in a vulnerability assessment.

To evaluate the most accurate assessment of extreme water levels, other drivers such as tides (e.g., King tides), seasonal and interannual effects (e.g., El Niño), river runoff from heavy precipitation, and shoreline characteristics of a given location, would also need to be accounted for in addition to storm conditions. This level of detail will likely not be necessary for all applications, but could be relevant for situations that are very risk averse to even a single flooding event (e.g., emergency services), or in landscapes that are especially vulnerable to the impacts of combined shoreline and watershed flooding (e.g., locations near bar-built estuaries).

BOX 15:

Storm Conditions

Coastal storms often bring high water levels along the coast. Contributors to coastal water levels include storm surge, wave runup, astronomical tides, seasonal variations, and regional sea level. These contributors to water levels are defined as follows:

- **Storm surge** is the temporary increase in sea level caused by low atmospheric pressure and wind during storms.
- **Wave runup** is the total rise in coastal water levels as waves break and rush up the beach.
- **Astronomical tides** are the regular rise and fall of the sea surface in response to forces exerted by the moon and sun.

- **Seasonal variations** in sea level occur when seasonal events cause temporary changes to sea levels. For example, El Niño brings warmer water that has a higher volume, resulting in increased water levels.

During coastal storms, especially those that occur at the same time as high astronomical tides and/or seasonal variations that cause increased water levels like El Niño events, coastal water levels can become extremely high. Extreme water levels, especially when coupled with land-based flooding from heavy precipitation, can jeopardize public safety and infrastructure. It is helpful for communities to have an understanding of how sea level rise may affect and exacerbate the severity of coastal storm impacts and increase their frequency.

Groundwater can be an additional contributor to flooding that should also be considered. A roughly 1:1 relationship between sea level rise and groundwater rise can extend as far as 0.5 miles inland from the shoreline. For example, if one foot of sea level rise is expected by 2050 at a particular location, groundwater may also rise one foot above its current level in areas adjacent to the shoreline.⁷³ For low lying areas, particularly bays and river valleys where groundwater levels are already near the surface, this can result in emergent flooding. Locations within three miles of the coast or shoreline should consider the potential impacts of rising groundwater, however, the 1:1 relationship will taper off further inland. Watershed-scale modeling is necessary to make specific local predictions, but in the absence of local modeling either the 1:1 assumption or the groundwater hazard projections in the OCOF map viewer may be used. In addition to potential contributions to flooding, rising groundwater can be a hazard of concern because it can mobilize subsurface contaminants that were previously above the water table, become salty as a result of pumping, and infiltrate or corrode underground infrastructure. For more information on groundwater resources and analysis see Section 3.2.

Ultimately, the project team, with input from relevant stakeholders, should choose sea level rise and storm scenarios for vulnerability assessments that are appropriate for a specific project, can be accomplished within the project budget, and are adequately precautionary. Depending on the type of vulnerability assessment pursued, evaluation of each additional scenario and/or storm condition may have an associated cost.

73. Befus et al., 2020.



BOX 16:

Coastal Storm Modeling System (CoSMoS) - Sea Level Rise Visualization Tool

CoSMoS is a dynamic modeling approach developed by the United States Geological Survey and accessed through the Our Coast, Our Future (OCOF) web interface.⁷⁴ As described in Box 12 above, OCOF is a user-friendly web-based tool for understanding and viewing potential flooding, coastal erosion, and rising groundwater hazards based on model outputs from CoSMoS. This tool allows users to identify areas along California's coastline that will likely be exposed to and are at risk from these coastal hazards due to projected sea level rise. Its flooding layer allows users to select different values of sea level rise (in meters and feet) and storm frequency (none [i.e., daily conditions], annual, 20-year, or 100-year) to visualize flooding at a 1-meter resolution across California's outer coast and estuaries (the North Coast is planned for release in 2024), including San Francisco Bay.

Because CoSMoS is a comprehensive modeling tool, it incorporates all the key drivers of coastal flooding, including sea level rise, tides, seasonal effects, storm surge, waves, and river flows to produce location specific maps showing total water levels and the associated flooding. It can be used both as a screening tool to examine the flooding from different combinations of sea level rise and storm conditions to help inform selection of the most appropriate sea level rise and storm scenarios for a vulnerability assessment (Step 3) and/or to support the exposure analysis of a vulnerability assessment (Step 4). For most planning and projects, OCOF is well-suited

to conduct the exposure analysis portion of a vulnerability assessment.

More specifically, to support screening and/or exposure analysis, users can:

- View locally relevant online maps and tools to understand vulnerabilities to sea level rise, storms, and shoreline change;
- Interact with a map that includes flood extent, depth, duration, wave heights, current velocity, cliff retreat, shoreline change, and groundwater emergence;
- View and download CoSMoS information through the OCOF⁷⁵ flood mapper, which provides a user-friendly web-based tool for viewing all model results;
- Download modeling results as GIS shapefiles with accompanying metadata at USGS ScienceBase-Catalog;⁷⁶ or
- View and interact with estimates of residents, businesses, and infrastructure that could be exposed to CoSMoS flooding projections from each coastal storm and sea level scenario through the Hazards Exposure Reporting and Analytics (HERA)⁷⁷ application.

OCOF allows users to select from a limited number of sea level values (i.e., 0.8, 1.6, 2.5, 3.3, 4.1, 4.9, 5.7, 6.6, 8.2, 9.8, and 16.4 feet). Depending on what sea levels are identified for analysis, there might not be an option that is an exact match. For instance, the Humboldt Bay Intermediate scenario at 2100 is 3.9 feet, which is between the available selections of 3.3 and 4.1 feet in OCOF. For these situations, it is not always necessary to analyze the exact numerical values; it is more important that the full range of values from Intermediate to High scenarios, with consideration of storm conditions, is evaluated.

74. <https://www.usgs.gov/centers/pcmssc/science/coastal-storm-modeling-system-cosmos#overview>

75. <http://ourcoastourfuture.org/>

76. <https://www.sciencebase.gov/catalog/item/5633fea2e4b048076347f1cf>

77. <https://www.usgs.gov/apps/hera/>

>> STEP 4: Conduct vulnerability assessment

A vulnerability assessment that evaluates impacts from potential sea level rise-induced inundation and flooding includes three key components: exposure, sensitivity, and adaptive capacity. Exposure is the degree to which habitats, people, private property, critical infrastructure, and public access will be affected by sea level rise. Sensitivity is the extent to which these natural and built assets will be damaged or destroyed by that exposure. And adaptive capacity is the ability of natural systems and infrastructure to respond or adapt to rising sea levels to minimize harm.

The first step in a vulnerability assessment is to create exposure maps of sea level rise induced inundation and flooding. This can be accomplished by using a sea level rise visualization tool, such as OCOF which provides visualizations of future flooding and inundation under different sea level rise projections using the CoSMoS model (see Box 16 for more detail on CoSMoS). However, the level of complexity, methodologies, and underlying assumptions differ across tools so it is important to understand these variations when selecting an appropriate visualization tool. Alternatively, a tailored made-to-order exposure mapping effort can be performed with more local considerations. These hyper-local and technologically advanced assessments can be time intensive and costly but can provide much greater detail and accuracy than the visualization tools available online. Additionally, aspects of sensitivity can be integrated into project-specific exposure mapping, for example coupling sea level rise exposure with habitat sensitivity information.

Erosion and groundwater are unique considerations because they can both contribute to exposure and are also systems that will be impacted. Erosion of beaches, cliffs, bluffs, dunes, and other landforms are important components of a vulnerability assessment and should be included to evaluate exposure and impacts. Similarly, groundwater can be integrated into an exposure analysis, however this requires local water table and geologic information that might not be readily attainable or might add significantly to the expense. For locations with shallow water tables, it is recommended that impacts to groundwater and saltwater intrusion be assessed since rising groundwater may pose a greater flood risk than tidal flooding in some areas.⁷⁸ Loss and destruction of other habitats/ecosystems (such as wetlands or submerged aquatic vegetation) can also contribute to exposure as well as being an impacted system. These ecosystem-level dynamics can be very difficult to capture unless an advanced tailor-made assessment is pursued.

Sensitivity is the degree to which a place or asset is affected by sea level rise inundation and temporary flooding. For flooding, this can be sensitivity to a one-time flood or most often, sensitivity to multiple flooding

78. Befus et al., 2020.

occurrences. For community planning, sensitivity (or impact) analysis should include flood frequency with sea level rise and evaluation of flood threshold (height and/or number of flood events), to understand how an asset or community is impacted. This should also include an assessment of impacts to environmental justice communities and consideration of erosion, coastal habitats and wildlife, saltwater intrusion and groundwater, recreational areas and access, energy infrastructure, transportation systems, flood protection infrastructure, wastewater systems, stormwater systems, community services and critical facilities, toxic sites, and landfills and waste facilities. For a site-specific project, the analysis would be limited and appropriate to the project scope and location.

The final step in a vulnerability assessment encourages the community to measure the degree to which it is equipped to adapt to sea level rise (i.e., adaptive capacity, see Box 17 below) through the existence of policies, structures, finances, and human resources that can assist, or already are assisting, adaptation to potential changes.

The project team, with input from relevant stakeholders, should conduct a vulnerability assessment that is appropriate for a specific project and can be accomplished within the project budget. It might also be appropriate to use a free online visualization tool to understand a broad range of risks at a high level, but only pursue a detailed analysis for the sea level value(s) most relevant to the project or the value(s) that would result in specific impacts of interest.

BOX 17:

Adaptive Capacity

Adaptive capacity is the ability of a system or community to evolve in response to, or cope with the impacts of sea level rise. Assets or natural resources with high adaptive capacity will likely have greater flexibility and potential to withstand rising sea levels. Adaptive capacity may be inherent to the asset or can be improved through forward-looking planning or design (e.g., including sufficient physical space to allow for buffering effects or inland migration of habitats, or designing a structure that can be easily relocated). Adaptive capacity

is also a function of the existing characteristics of a system. For example, a community that is chronically under-resourced may develop effective adaptation strategies, but will likely still be at a disadvantage compared to communities with more resources for advanced planning and implementation. This lack of adaptive capacity is a barrier to equitable adaptation.

See Chapter 4.3 for additional recommendations for sea level rise planning and adaptation, including information on prioritizing social equity, environmental justice, and the needs of underserved and vulnerable communities.

>> STEP 5: Explore adaptation options and feasibility

The results of the vulnerability assessment should highlight what is most vulnerable and allow identification of adaptation priorities. This should be done in close coordination with the community, stakeholders, and relevant regulatory bodies. Typically, the next step is to explore site-specific adaptation options and the feasibility of these options, either through an adaptation pathways approach or as a standalone project. This could include discussion of all available adaptation strategies and selection of a limited number that are brought to a conceptual design stage or conducting a formal alternatives analysis. A cost-benefit analysis could also be conducted at this stage.

Often the suite of adaptation options is limited by legal, physical, economic, or social constraints. Evaluating adaptation options will be a highly local process that includes the values and priorities of a particular community.

>> STEP 6: Select phased adaptation approach and/or implement project

Following a thorough assessment of adaptation options, a specific project or adaptation pathway must ultimately be selected. The process of selecting an implementation project or adaptation pathway (i.e., a project design along with future, trigger-based adaptation actions) that is adaptive to a certain amount of sea level rise will include consideration of risk, budget, regulatory constraints, environmental and community impacts, and stakeholder input, in addition to other factors. Selection will often be a negotiation and assessment of trade-offs, and using trigger-based adaptation pathways to account for sea level rise over time - rather than incorporating the full sea level rise amount into initial project design - will likely be a common approach.



BOX 18:**How to Evaluate Risk for a Project**

Risk aversion is the strong inclination to avoid taking risks in the face of uncertainty. State and local governments should consider the risks associated with various Sea Level Scenarios and determine their tolerance for, or aversion to, those risks (i.e., over or underestimating the amount of sea level rise). Risk aversion and development type are not strictly defined boxes but rather represent a spectrum of risk levels that in practice need to be evaluated on a project-by-project basis. For many situations, there will be multiple development types within a single project or planning effort that will need to be considered. Additionally, the lifespan of a project or adaptation action, is a major factor in its risk profile. There is no quantitative calculation to determine risk level, however general guidance can be provided:

LOW RISK AVERSION:

Adaptative, short lifespan (i.e., 2050-2075), lower consequence situations that are fairly risk tolerant, for instance a public bench or coastal recreational trail. Additionally, low risk should be considered for managing or restoring natural infrastructure, such as tidal wetland management and restoration, creating living shorelines, estimation of saltwater intrusion and coastal landscape migration, or protecting estuarine water quality. When an action has the capacity to be adapted or is not anticipated to persist long into the future, it is often more important to know the most likely sea level rise rather than the maximum plausible sea level rise. Therefore, if an adaptation action is part of a short lifespan adaptation pathway, a development type that might otherwise fall into a higher risk aversion category could be considered low risk in this situation. Low risk aversion projects should be resilient to the Intermediate Scenario.

MEDIUM-HIGH RISK AVERSION:

Appropriate for less adaptive, long lifespan (i.e., 2075 and beyond), more vulnerable projects that will experience medium to high consequences if impacted because of underestimating sea level rise. Multi-use paths that provide public access and/or are part of a transportation network may fall into the

medium-high risk aversion category. Most commercial and residential development will fall in the medium-high or extreme risk averse categories, depending on the size and scale of the development. These efforts should be resilient to the Intermediate-High Scenario, when feasible.

EXTREME RISK AVERSION:

For high consequence and very long lifespan (i.e., beyond 2100) projects that have little to no adaptive capacity, would be irreversibly destroyed or significantly costly to relocate/repair and would have considerable public health, public safety, or environmental impacts. For instance, critical infrastructure which includes but is not limited to, transit, roads, airports, ports, water storage and conveyance, wastewater treatment facilities, landfills, power plants, and railroads,) should be considered as extremely risk averse. Extreme risk aversion projects should be resilient to the High Scenario, through project design or through an adaptation pathway approach, when feasible and appropriate.

For short-term adaptation actions (e.g., as part of an adaptation pathways approach) the Intermediate Scenario is recommended, regardless of risk category. This is because multiple lines of evidence identify the Intermediate Scenario as being most likely in the near-term (i.e. 2050).

For longer-term actions, general guidance can be provided, from a risk perspective, on how to select a Sea Level Scenario. For low-risk averse projects, it is recommended that the Intermediate Scenario be applied, which has a 5% chance of being exceeded in 2100 (assuming 3°C of warming and no low confidence ice sheet processes). Because there is an 82% chance that the Intermediate-Low Scenario will be exceeded in 2100 (assuming 3°C of warming and no low confidence ice sheet processes), the Intermediate-Low Scenario is not recommended even for low-risk averse projects.

For medium-high risk averse applications, the Intermediate-High Scenario is recommended. Although there is a 0.1% chance of exceeding the Intermediate-High Scenario in 2100 (assuming 3°C of warming and no low confidence processes) the state recommends a precautionary approach, when possible, to maximize preparedness and resilience. Furthermore, if there is greater than 4°C warming and contribution from low confidence processes, there is a 20% chance of exceeding the Intermediate-High Scenario in 2100 (high levels of warming). Additionally, because medium-risk averse projects have longer lifespans, the Intermediate-High Scenario provides an additional buffer should the project need to persist further into the future than originally planned for.

For extreme risk averse applications, the High Scenario may be appropriate, however, there are limited situations in which designing and constructing to the High Scenario will be necessary and/or feasible without significant logistical tradeoffs. If significant constraints do not exist, then designing to the High Scenario is recommended, all other factors being equal. However, it is likely that in most

situations, factors like the urbanized nature of existing communities, location of existing facilities, requirements to provide service to existing development, and fiscal constraints will make incorporating the High Scenario into initial project design infeasible. The adaptation pathways approach is therefore recommended, in which a smaller amount of sea level rise is incorporated into initial project design while also developing options to address higher sea level rise amounts in the future. This may include high level planning for the types of strategies that may be necessary if and when the amounts of sea level rise represented in the High Scenario occurs, identification of triggers for additional adaptation planning, or development of adaptation pathways that include triggers for additional adaptation measures or redesign to implement if certain thresholds of impacts or sea level rise amounts occur.

At a minimum, the High Scenario should be analyzed (Steps 3 and 4) to understand the associated impacts and to broadly understand what types of strategies may be needed, and the possibility of the High Scenario should be disclosed and monitored. Although the High Scenario has an effectively zero percent chance of being exceeded in 2100 (assuming 3°C of warming and no low confidence processes), extreme risk averse projects have anticipated lifespans beyond 2100 and therefore should be prepared for both worst case values at 2100, as well as higher amounts of sea level rise that are expected beyond 2100. Additionally, assuming high levels of warming in 2100 and contributions from the low confidence processes, there is an 8% chance of exceeding the High Scenario in 2100; therefore, it is still recommended to consider as part of worst-case planning.

Depending on the results of the vulnerability assessment, projects or plans might also need to account for extreme water levels that will occur due to storms, and/or rising groundwater. This should be assessed on a case-by-case basis and may result in planning or designing projects to a higher water level than a given Sea Level Scenario.

However, in a real-world situation, there might be a maximum level of sea level rise that can be incorporated into a project. Rather than selecting a sea level rise value to design to, a project's design is evaluated to assess if it is sufficiently protective of sea level rise and storm conditions over the lifespan of the project. It is important to maintain a transparent process, guided by the scientific recommendations, when making a final selection.

4.3. General Recommendations for Sea Level Rise Planning and Adaptation

The 2018 California Sea-Level Rise Guidance provided a set of recommendations to encourage sea level rise planning in alignment with state policy goals and priorities. These recommendations are carried forward in this 2024 Guidance Update below and provide guidance on preferred sea level rise planning and adaptation approaches, with an understanding that the diversity of communities, uses, and natural resources along California's coastline, as well as planning for new development versus existing structures, may merit different approaches to building resilience.

1. Adaptation planning and strategies should prioritize social equity, environmental justice and the needs of underserved and vulnerable communities.

Communities of color, low-income communities, and California Native American tribes have been, and will continue to be, disproportionately overburdened by pollution and climate change. Sea level rise will add to those burdens. Impacts such as increased flooding, damage to homes and roads, disruption to public transportation, elevated exposure to toxic materials, and destruction of coastal sacred places and cultural sites will unduly affect vulnerable communities. These impacts can manifest as complete community displacement, loss of areas with cultural and/or

historic significance, loss of personal property, worsened health, reduced or lost wages, and loss of free or affordable public access to the coast. Vulnerable communities may lack financial or other resources to plan for sea level rise as well as the ability to adequately respond to impacts once they occur.

Sea level rise planning that prioritizes social equity, environmental justice and protection of the lives and property of underserved and vulnerable communities should include early public engagement of those who will be directly or indirectly affected by rising sea levels, a focused characterization of impacts on exposed populations and communities dependent on critical assets threatened by sea level rise, and identification of specific adaptation strategies to minimize or mitigate these impacts. Engaging communities that face existing inequalities already (or will face unequal distribution of sea level rise impacts) early in the planning process will ensure that vulnerability assessments and adaptation strategies accurately reflect their risk, needs and priorities. State and local governments should also prioritize technical support and funding opportunities for planning and adaptation efforts of vulnerable, underserved, and tribal communities. Incorporating social equity and environmental justice in sea level rise planning and adaptation strategies should:

- **Address environmental contamination risks for coastal communities adjacent to industrial or toxic sites.** Coastal environmental justice communities tend to have fewer beachfront homes at risk of inundation but are often separated from the coast by strips of industrial facilities, ports and military installations. Sea level rise threatens job sites for local residents, risks mobilizing contamination from cleanup sites, and can damage critical energy, transportation or other infrastructure. Prioritizing cleanup of sites threatened by sea level rise and rising groundwater can prevent toxic contamination from migrating into nearby communities.

- **Preserve access to and along beaches, coastal wetlands, and other natural shoreline features.** Protecting natural coastlines preserves affordable outdoor recreation access for communities that often lack parks or other sources of green space and face existing health disparities. While many coastal cities in California include expensive beachfront homes, the coast is used regularly for recreation by working-class residents who are visiting or live nearby. Sea level rise planning and adaptation strategies should protect public access to and along the beach to maximize free or affordable use of the coast for the benefit of all people.
 - **Prevent displacement by ensuring that investments in coastal resilience protect local jobs and housing costs.** In climate adaptation policies, it is important to understand the economic ties between vulnerable communities and polluting industries along their coasts, and how to build environmentally healthy and economically vibrant communities. Deindustrialization of coastal areas and restoration of natural coastal habitats can result in major environmental benefits, but also job losses and rent increases for the very same communities who are intended to be protected by these natural buffers. Coastal resilience investments should provide economic benefits for adjacent working-class communities, including anti-displacement housing policies and local jobs programs.
 - **Address economic impacts on agriculture.** California has major agricultural regions along the Central Coast - such as the Oxnard Plain, Santa Maria Valley and Salinas Valley where tens of thousands of farmworkers are employed in the fields and whose livelihoods are threatened by seawater intrusion into groundwater aquifers. Focused monitoring of seawater intrusion in coastal agricultural areas, restoration of coastal wetlands buffers, and effective groundwater management to prevent excessive pumping and restore fresh groundwater could help prevent major long-term economic damage to agriculture and farmworkers.
 - **Address emergency response to and recovery from natural disasters.** Vulnerable populations including low-income, unhoused, elderly, disabled and immigrant communities, are often left behind in access to information and resources in the chaos of disaster response. Proactive, deliberate planning in partnership with marginalized communities can prepare for a major flooding disaster and minimize or mitigate the likelihood of systemic failure. Emergency services agencies should be prepared to coordinate with local, regional and state partners, including Listos California, on development and dissemination of multilingual communications that fulfill the language and access needs of local vulnerable communities. Consistent with the State Hazard Mitigation Plan, local emergency management plans should undergo regular review, and consider the effects of climate change, and the potential disproportionate impact incidents may have on vulnerable communities.
 - **Evaluate the social and economic implications of various adaptation strategies.** Planning and investment decisions that will increase risk to vulnerable communities should be avoided, and actions to bolster resilience and social equity should be prioritized.
- 2. Adaptation strategies should prioritize protection of coastal habitats and public access.**
- **Implement natural solutions for shoreline protection.** Strategies to protect shoreline development from sea level rise impacts should prioritize the use of nature-based solutions where feasible or appropriate and minimize shoreline armoring and flood barriers where possible. While hard structures or gray solutions provide

temporary protection against the threat of sea level rise, they disrupt natural shoreline processes, accelerate long-term erosion, and can prevent coastal habitats from migrating with sea level rise, causing loss of beaches and other critical habitats that provide ecosystem benefits for both wildlife and people; therefore, they should only be used in appropriate locations and situations. Hybrid projects, which combine nature-based approaches with traditional engineering “grey” approaches (for example, pocket beaches between artificial headlands), may be effective where purely nature-based approaches are not likely to meet design goals.

Natural shoreline infrastructure means utilizing the natural function of ecological systems or processes to reduce vulnerability to specific environmental hazards and increase resilience of the shoreline in order to perpetuate or restore its ecosystem services.⁷⁹ Natural infrastructure includes preservation or restoration of dunes, wetlands and other coastal habitats and leverages natural processes to reduce risk to human lives, property and infrastructure by providing a buffer against storm surge and increased wave action, thus reducing shoreline impacts and coastal erosion. These solutions have been shown in many cases to be low maintenance, cost-effective and adaptive to changing conditions. Additionally, natural infrastructure provides multiple benefits beyond flood protection including public access, habitat for wildlife and improved water quality, thereby building resilience while improving the overall ecological function of coastal systems.

In addition to prioritizing natural infrastructure, strategic relocation should be considered as a possible adaptation strategy to address rising sea levels. This can result in a landward redevelopment pattern and thoughtful realignment of development along the coast so that natural erosion and other coastal

processes, including beach formation, can continue. This approach also allows shorelines to migrate naturally, rather than using seawalls, flood barriers, or rock revetments to anchor them in a specific location and may involve changing patterns of residential, commercial, or industrial development and restoration of natural areas to enhance ecosystem services, make sound infrastructure investments, and provide additional protection and safety against flooding through buffering effects, as described above.

Strategic relocation will also provide added protection for wetlands, marshes and other important coastal habitats that will face inundation or erosion if restricted from moving landward by existing development or shoreline armoring. Decision makers should prioritize conservation, restoration and land acquisition of properties that can provide needed open space to accommodate inland migration in order to preserve the natural function of wetlands and other coastal ecosystems.

Restoration of wetlands and other coastal habitats should remain a priority in California even in the face of rising seas; even if present-day restored wetlands transition to subtidal habitat sometime in the future, there will still be continued ecosystem benefits for wildlife and people over the long term. In addition, wetland restoration and other adaptation strategies that provide greenhouse gas reduction benefits by storing and sequestering carbon should be prioritized.

- ***Preserve public access, including beaches and coastal parks, while protecting natural resources.*** Public access along California’s coast is already being affected by sea level rise, coastal flooding, and erosion. Coastal trails, public beaches, park infrastructure,

79. Newkirk et al, 2018.

and other state and public assets that are of high value to Californians will increasingly be under threat from higher sea levels, intensified wave action, and accelerated coastal erosion.

Decision makers, including state and local agencies that manage state or locally owned coastal assets, should assess the vulnerability of public access and prioritize its protection for the invaluable benefits it provides to residents and visitors. Every effort should be made to ensure that protection of public access or park infrastructure does not degrade coastal habitats. Beaches backed by development or shoreline armoring will not be able to migrate inland as sea levels rise, resulting in permanent inundation over time and loss of public access. Consideration should be given to allowing for natural shoreline movement and relocation of public access and park infrastructure to preserve beach access and protect wetlands, dunes and other coastal habitats. Using natural infrastructure to safeguard public access facilities, parks, and trails or planning ahead to relocate these resources will help ensure that both public access and coastal habitats are preserved for the long-term.

3. Adaptation strategies should consider the unique characteristics, constraints and values of existing water-dependent infrastructure, ports and Public Trust uses.

Existing water-dependent infrastructure and ports support Public Trust uses vital to the State (such as commerce, navigation, fisheries, and recreation) and have unique characteristics and constraints for adaptation to sea level rise. They are often located in densely developed coastal areas where asset relocation, natural infrastructure solutions, and other space-dependent strategies may not be feasible. Planners should continue to collaborate regionally and with the State to develop adaptation strategies for water-dependent infrastructure that will be protected

in place, as well as address strategies to adapt existing infrastructure into the future. Existing shoreline protective structures may need to be repaired and retrofitted to adapt to rising sea levels. Negative impacts to other Public Trust values, including coastal habitats and public access, should be minimized in all existing and future use of shoreline protective structures. Innovative and resilient design alternatives to conventional gray infrastructure should be explored when retrofitting existing protective structures or contemplating future protective structures.

4. Consider episodic increases in sea level rise caused by storms and other extreme events.

As described in more detail in Chapter 3.0 above, individually or in combination, these events will produce significantly higher water levels than sea level rise alone, and will likely be the drivers of the strongest impacts to coastal ecosystems, development and public access over the next several decades. Water levels reached during these large, acute events have already caused significant damage along California's coast. For example, a strong El Niño combined with a series of storms during high-tide events caused more than \$200 million in damage (in 2010 dollars) to the California coast during the winter of 1982-83. Additionally, in areas where rivers meet the ocean, the combined effects of sea level rise, storm conditions and higher riverine water levels could further exacerbate flooding conditions in these locations.

Furthermore, climate change may result in increased frequency or intensity of coastal storms and extreme events, posing even greater risks for California's coastline from flooding, erosion and wave damage. To adequately protect coastal communities, infrastructure and natural resources, decision makers should consider extreme oceanographic conditions in conjunction with sea level rise over the expected life of a project. A range of existing mapping tools is

available to help evaluate storm-related coastal flooding, sea-level rise and shoreline change and to evaluate impacts and change into the future; these mapping tools are described in detail below. In addition to these tools, the San Francisco Bay Conservation and Development Commission's (BCDC) Adapting to Rising Tides (ART) Program has developed robust and locally-relevant resources for the San Francisco Bay to understand current and future flood risk.⁸⁰ It is important to note that current Federal Emergency Management Agency (FEMA) flood maps are based on existing shoreline characteristics and wave and storm climatology at the time of the flood study and historic storm data; therefore, these maps will not reflect flood hazards based on anticipated future sea levels or increased storms associated with climate change.⁸¹

5. Coordinate and collaborate with local, state and federal agencies when selecting sea level rise scenarios; where feasible, use consistent sea-level rise scenarios across multi-agency planning and regulatory decisions.

Project planning and design along the coast often requires approval by multiple agencies across local, regional, state and federal levels. To increase efficiency and standardize risk evaluation, efforts led by or under the regulatory authority of multiple agencies should use the same sea level rise scenarios to achieve consistency across specific projects and regions. Cross-jurisdictional decisions should also prioritize implementation of consistent or complementary adaptation strategies. Some tools already exist to support these types of cross-jurisdictional adaptation planning decisions within regions, including the San Francisco Bay Shoreline Adaptation Atlas.⁸²

6. Consider local conditions to inform decision making.

Local circumstances and associated sea level rise impacts should be assessed to inform adaptation decisions that will protect communities and the environment. The

interplay between sea level rise and conditions such as contaminated soil, groundwater, or stormwater systems as well as beach and cliff erosion can vary significantly along the coast and should be evaluated at a local level. The diversity of shoreline types, natural conditions, community characteristics, services, assets, land ownership, and local priorities may warrant different approaches to planning and adaptation, particularly when making decisions for new development versus maintenance or replacement of existing assets necessary for public health and safety. Adaptation pathways with a phased approach can invoke the precautionary principle while maintaining protection of community well-being, the environment, and critical assets.

7. Assessment of risk and adaptation planning should be conducted at community and regional levels, when possible.

Sea level rise planning decisions made for one municipality, or even one landowner, have the potential to impact the resiliency of nearby properties and coastal habitats. A jurisdiction that chooses to implement natural infrastructure may lose some of the benefits and protection from this adaptation strategy if an adjacent community decides to construct a seawall. Decision makers should identify opportunities to coordinate regional adaptation planning efforts by: conducting regional vulnerability assessments to evaluate common risks; leveraging technical and financial resources; and implementing consistent regional adaptation strategies. BCDC's ART Program and the San Diego Regional Climate Collaborative⁸³ are examples of regional planning efforts that can serve as models for other regional planning efforts throughout the state.

80. www.adaptingtorisingtides.org

81. <https://www.fema.gov/flood-maps/coastal#How>

82. Beagle et al., 2019.

83. <https://www.sdclimatecollaborative.org>

5. Conclusions and Looking Forward

AUTHORS: Ocean Protection Council and Ocean Science Trust

CALIFORNIA MUST take bold and swift action to protect nature and coastal communities from the impacts of sea level rise. This action needs to be grounded in current science, standardized across jurisdictions, and flexible enough to accommodate local priorities while ensuring that the state is adequately prepared to adapt to the expected changes ahead.

As demonstrated in this update, understanding of sea level rise continues to rapidly evolve with increases in data availability and scientific tools. The IPCC, along with U.S. federal agencies, is expected to provide updates on sea level rise trends and projections every five years, which will continue to serve as the foundation for updates to California's sea level rise guidance. Over the next five years, we anticipate that scientific understanding will be further refined, leading to even more precise guidance on anticipated sea level rise and the use of scenarios for adaptation planning and projects. Monitoring sea level rise trends and impacts for adaptive management and planning will continue to be a key part of this process.

This report, based on science from a team of leading experts, provides overarching science and policy guidance to support coordinated and consistent planning and adaptation. Affording local decision-making autonomy is important to ensure that planning is location-specific and tailored to circumstances – and is guided by overarching science. This guidance allows for local planners, elected officials, tribes and additional decisionmakers to make the most appropriate decisions for their communities. OPC is committed to continuing to provide scientific support and build capacity for sea level rise planning and adaptation. OPC actions moving forward include:

- Ongoing coordination with the existing membership of the State Sea Level

Rise Collaborative, including continued implementation, progress accounting, and future update of the State Agency Sea Level Rise Action Plan for California.

- Prioritized integration of local, regional, and tribal governments on the State Sea Level Rise Collaborative to further embed local priorities and needs into statewide policies and actions.
- Accelerated access to funding for standardized sea level rise adaptation plans and projects through OPC's Senate Bill (SB) 1 Grant Program; maximized investments to highly vulnerable and under-resourced communities.
- Partnership with the Governor's Office of Planning and Research, California Natural Resources Agency, the California Energy Commission, and the Strategic Growth Council to bridge this guidance with the sea level rise research and recommendations included in California's Fifth Climate Change Assessment scheduled for completion in 2026.
- Integration of updated sea level rise scenarios and policy guidance into OPC's broader Strategic Plan priorities to build climate resilience for marine ecosystems and communities, advance equity, conserve biodiversity, and promote the sustainable blue economy.

California can continue to serve as a model for the nation for how to integrate science into policy and planning decisions to minimize impacts from sea level rise on ecosystems, cultural resources and traditional practices, livelihoods, and public and private property. There is much work ahead, but collective action at the local, regional, tribal, state and federal levels will build the resilience that California needs to thrive, even in a changing climate.

6. References

- Adapting Stormwater Management for Coastal Floods. (n.d.). Retrieved August 9, 2023, from <https://coast.noaa.gov/stormwater-floods/>
- Adapting to Rising Tides. (n.d.). Retrieved August 9, 2023, from <https://www.adaptingtorisingtides.org/>
- Aerts, R., Honnay, O., & Van Nieuwenhuysse, A. (2018). Biodiversity and human health: Mechanisms and evidence of the positive health effects of diversity in nature and green spaces. *British Medical Bulletin*, 127(1), 5–22. <https://doi.org/10.1093/bmb/ldy021>
- Apg | ResilientCA. (n.d.). Retrieved August 8, 2023, from <https://resilientca.org/apg/>
- ART Bay Shoreline Flood Explorer. (n.d.). Retrieved August 9, 2023, from <https://explorer.adaptingtorisingtides.org/explorer>
- Barbier, E. B. (2013). Valuing Ecosystem Services for Coastal Wetland Protection and Restoration: Progress and Challenges. *Resources*, 2(3), 213–230. <https://doi.org/10.3390/resources2030213>
- Barnard, P. L., Dugan, J. E., Page, H. M., Wood, N. J., Hart, J. A. F., Cayan, D. R., Erikson, L. H., Hubbard, D. M., Myers, M. R., Melack, J. M., & Iacobellis, S. F. (2021). Multiple climate change-driven tipping points for coastal systems. *Scientific Reports*, 11(1), 15560. <https://doi.org/10.1038/s41598-021-94942-7>
- Barnard, P. L., Erikson, L. H., Foxgrover, A. C., Hart, J. A. F., Limber, P., O'Neill, A. C., van Ormondt, M., Vitousek, S., Wood, N., Hayden, M. K., & Jones, J. M. (2019). Dynamic flood modeling essential to assess the coastal impacts of climate change. *Scientific Reports*, 9(1), 4309. <https://doi.org/10.1038/s41598-019-40742-z>
- Barnard, P. L., Hoover, D., Hubbard, D. M., Snyder, A., Ludka, B. C., Allan, J., Kaminsky, G. M., Ruggiero, P., Gallien, T. W., Gabel, L., McCandless, D., Weiner, H. M., Cohn, N., Anderson, D. L., & Serafin, K. A. (2017). Extreme oceanographic forcing and coastal response due to the 2015–2016 El Niño. *Nature Communications*, 8(1), 14365. <https://doi.org/10.1038/ncomms14365>
- Barnard, P. L., Short, A. D., Harley, M. D., Splinter, K. D., Vitousek, S., Turner, I. L., Allan, J., Banno, M., Bryan, K. R., Doria, A., Hansen, J. E., Kato, S., Kuriyama, Y., Randall-Goodwin, E., Ruggiero, P., Walker, I. J., & Heathfield, D. K. (2015). Coastal vulnerability across the Pacific dominated by El Niño/Southern Oscillation. *Nature Geoscience*, 8(10), 801–807. <https://doi.org/10.1038/ngeo2539>
- Barnett, J., Graham, S., Mortreux, C., Fincher, R., Waters, E., & Hurlimann, A. (2014). A local coastal adaptation pathway. *Nature Climate Change*, 4(12), 1103–1108. <https://doi.org/10.1038/nclimate2383>
- Beagle, J., Lowe, J., McKnight, K., Safran, S. M., Tam, L., & Szambelan, S. Jo. (2019). San Francisco Bay Shoreline Adaptation Atlas: Working with Nature to Plan for Sea Level Rise Using Operational Landscape Units. SFEI Contribution No. 915. SFEI & SPUR: Richmond, CA. p 255. <https://www.sfei.org/documents/adaptationatlas>
- Befus, K. M., Barnard, P. L., Hoover, D. J., Finzi Hart, J. A., & Voss, C. I. (2020). Increasing threat of coastal groundwater hazards from sea level rise in California. *Nature Climate Change*, 10(10), 946–952. <https://doi.org/10.1038/s41558-020-0874-1>

- Bekaert, D. P. S., Hamlington, B. D., Buzzanga, B., & Jones, C. E. (2017). Spaceborne Synthetic Aperture Radar Survey of Subsidence in Hampton Roads, Virginia (USA). *Scientific Reports*, 7(1), 14752. <https://doi.org/10.1038/s41598-017-15309-5>
- Bekaert, D. P. S., Jones, C. E., An, K., & Huang, M.-H. (2019). Exploiting UAVSAR for a comprehensive analysis of subsidence in the Sacramento Delta. *Remote Sensing of Environment*, 220, 124-134. <https://doi.org/10.1016/j.rse.2018.10.023>
- Box, J. E., Hubbard, A., Bahr, D. B., Colgan, W. T., Fettweis, X., Mankoff, K. D., Wehrlé, A., Noël, B., van den Broeke, M. R., Wouters, B., Bjørk, A. A., & Fausto, R. S. (2022a). Greenland ice sheet climate disequilibrium and committed sea level rise. *Nature Climate Change*, 12(9), 808-813. <https://doi.org/10.1038/s41558-022-01441-2>
- Box, J. E., Hubbard, A., Bahr, D. B., Colgan, W. T., Fettweis, X., Mankoff, K. D., Wehrlé, A., Noël, B., van den Broeke, M. R., Wouters, B., Bjørk, A. A., & Fausto, R. S. (2022b). Greenland ice sheet climate disequilibrium and committed sea level rise. *Nature Climate Change*, 12(9), 808-813. <https://doi.org/10.1038/s41558-022-01441-2>
- Bromirski, P. D. (2023). Climate-Induced Decadal Ocean Wave Height Variability From Microseisms: 1931-2021. *Journal of Geophysical Research: Oceans*, 128(8), e2023JC019722. <https://doi.org/10.1029/2023JC019722>
- Bromirski, P. D., & Kossin, J. P. (2008). Increasing hurricane wave power along the U.S. Atlantic and Gulf coasts. *Journal of Geophysical Research: Oceans*, 113(C7). <https://doi.org/10.1029/2007JC004706>
- Bromirski, P. D., Miller, A. J., Flick, R. E., & Auad, G. (2011). Dynamical suppression of sea level rise along the Pacific coast of North America: Indications for imminent acceleration. *Journal of Geophysical Research: Oceans*, 116(C7). <https://doi.org/10.1029/2010JC006759>
- Buffington, K. J., Janousek, C. N., Dugger, B. D., Callaway, J. C., Schile-Beers, L. M., Sloane, E. B., & Thorne, K. M. (2021). Incorporation of uncertainty to improve projections of tidal wetland elevation and carbon accumulation with sea level rise. *PLOS ONE*, 16(10), e0256707. <https://doi.org/10.1371/journal.pone.0256707>
- Buzzanga, B., Bekaert, D. P. S., Hamlington, B. D., & Sangha, S. S. (2020). Toward Sustained Monitoring of Subsidence at the Coast Using InSAR and GPS: An Application in Hampton Roads, Virginia. *Geophysical Research Letters*, 47(18), e2020GL090013. <https://doi.org/10.1029/2020GL090013>
- Cai, W., Borlace, S., Lengaigne, M., van Rensch, P., Collins, M., Vecchi, G., Timmermann, A., Santoso, A., McPhaden, M. J., Wu, L., England, M. H., Wang, G., Guilyardi, E., & Jin, F.-F. (2014). Increasing frequency of extreme El Niño events due to greenhouse warming. *Nature Climate Change*, 4(2), 111-116. <https://doi.org/10.1038/nclimate2100>
- Cai, W., Ng, B., Wang, G., Santoso, A., Wu, L., & Yang, K. (2022). Increased ENSO sea surface temperature variability under four IPCC emission scenarios. *Nature Climate Change*, 12(3), 228-231. <https://doi.org/10.1038/s41558-022-01282-z>
- Cal-Adapt. (n.d.). Retrieved August 9, 2023, from <https://cal-adapt.org/>
- California, S. of. (n.d.). SB 1 Funding. California Ocean Protection Council. Retrieved April 30, 2024 from <https://opc.ca.gov/sb-1-funding/>
- California Coastal Commission. (2023). Public Trust Guiding Principles and Action Plan. https://documents.coastal.ca.gov/assets/public-trust/Public%20Trust%20Guidance%20and%20Action%20Plan_Adopted.pdf

- California Governor’s Office of Planning and Research (OPR). (2017). Planning and Investing for a Resilient California: A Guidebook for State Agencies. California Governor’s Office of Planning and Research. <https://resilientca.org/projects/aafbf831-a4f0-47a6-8064-c6009a2f2c35/>
- California Natural Resources Agency. (2020). Making California’s Coast Resilient to Sea Level Rise: Principles for Aligned State Action. California Natural Resources Agency. http://www.opc.ca.gov/webmaster/_media_library/2020/05/State-SLR-Principles_FINAL_April-2020.pdf
- California State Lands Commission. (2023). Shoreline Adaptation and the Public Trust: Protecting California’s Public Trust Resources from Sea Level Rise. State of California. <https://slcprdwordpressstorage.blob.core.windows.net/wordpressdata/2023/12/Shoreline-Adaptation-Report.pdf>
- Callahan, C. W., Chen, C., Rugenstein, M., Bloch-Johnson, J., Yang, S., & Moyer, E. J. (2021). Robust decrease in El Niño/Southern Oscillation amplitude under long-term warming. *Nature Climate Change*, 11(9), 752–757. <https://doi.org/10.1038/s41558-021-01099-2>
- Cherry, J. A., McKee, K. L., & Grace, J. B. (2009). Elevated CO₂ enhances biological contributions to elevation change in coastal wetlands by offsetting stressors associated with sea level rise. *Journal of Ecology*, 97(1), 67–77. <https://doi.org/10.1111/j.1365-2745.2008.01449.x>
- Christensen, J., & King, P. (2017). Access for All. A New Generation’s Challenges on the California Coast. Institute of the Environment & Sustainability. University of California, Los Angeles. Los Angeles, California. <https://www.ioes.ucla.edu/wp-content/uploads/2017/01/UCLA-Coastal-Access-Policy-Report.pdf>
- Coastal Flood Risk | FEMA.gov. (2021, March 22). <https://www.fema.gov/flood-maps/coastal>
- Coastal Resilience – Mapping portal. (n.d.). Retrieved April 30, 2024, from <https://maps.coastalresilience.org/>
- Coastal Storm Modeling System (CoSMoS) | U.S. Geological Survey. (n.d.). Retrieved August 9, 2023, from <https://www.usgs.gov/centers/pcmsc/science/coastal-storm-modeling-system-cosmos>
- Collini, R. C., Heming, M. C., Mohrman, C., Daigle, M. T., Fulford, C. A., Lowry, C. L. G., Hanisko, M. D., Mikulencak, S., Price, R., Ransom, K. R., Sempier, T. T., Shepard, C., Underwood, W. V., Woodrey, M. S., Denny, M. D., & Sparks, E. (2022). Utilizing an End-User Driven Process to Identify and Address Climate-Resilience Tool Needs in the U.S. Gulf of Mexico. *Coastal Management*, 50(2), 197–214. <https://doi.org/10.1080/08920753.2022.2022975>
- Collins, M., An, S.-I., Cai, W., Ganachaud, A., Guilyardi, E., Jin, F.-F., Jochum, M., Lengaigne, M., Power, S., Timmermann, A., Vecchi, G., & Wittenberg, A. (2010). The impact of global warming on the tropical Pacific Ocean and El Niño. *Nature Geoscience*, 3(6), 391–397. <https://doi.org/10.1038/ngeo868>
- Dangendorf, S., Hay, C., Calafat, F. M., Marcos, M., Piecuch, C. G., Berk, K., & Jensen, J. (2019). Persistent acceleration in global sea level rise since the 1960s. *Nature Climate Change*, 9(9), 705–710. <https://doi.org/10.1038/s41558-019-0531-8>
- de Ruig, L. T., Barnard, P. L., Botzen, W. J. W., Grifman, P., Hart, J. F., de Moel, H., Sadrpour, N., & Aerts, J. C. J. H. (2019). An economic evaluation of adaptation pathways in coastal mega cities: An illustration for Los Angeles. *Science of The Total Environment*, 678, 647–659. <https://doi.org/10.1016/j.scitotenv.2019.04.308>

- DeConto, R. M., & Pollard, D. (2016). Contribution of Antarctica to past and future sea level rise. *Nature*, 531(7596), 591–597. <https://doi.org/10.1038/nature17145>
- DeConto, R. M., Pollard, D., Alley, R. B., Velicogna, I., Gasson, E., Gomez, N., Sadai, S., Condrón, A., Gilford, D. M., Ashe, E. L., Kopp, R. E., Li, D., & Dutton, A. (2021). The Paris Climate Agreement and future sea level rise from Antarctica. *Nature*, 593(7857), 83–89. <https://doi.org/10.1038/s41586-021-03427-0>
- Dugan, J. E., & Hubbard, D. M. (2010). Loss of Coastal Strand Habitat in Southern California: The Role of Beach Grooming. *Estuaries and Coasts*, 33(1), 67–77. <https://doi.org/10.1007/s12237-009-9239-8>
- Dugan, J. E., Hubbard, D. M., Rodil, I. F., Revell, D. L., & Schroeter, S. (2008). Ecological effects of coastal armoring on sandy beaches. *Marine Ecology*, 29(s1), 160–170. <https://doi.org/10.1111/j.1439-0485.2008.00231.x>
- Erikson, L. H., Hegermiller, C. A., Barnard, P. L., Ruggiero, P., & van Ormondt, M. (2015). Projected wave conditions in the Eastern North Pacific under the influence of two CMIP5 climate scenarios. *Ocean Modelling*, 96, 171–185. <https://doi.org/10.1016/j.ocemod.2015.07.004>
- Erikson, L., Morim, J., Hemer, M., Young, I., Wang, X. L., Mentaschi, L., Mori, N., Semedo, A., Stopa, J., Grigorieva, V., Gulev, S., Aarnes, O., Bidlot, J.-R., Breivik, Ø., Bricheno, L., Shimura, T., Menendez, M., Markina, M., Sharmar, V., ... Webb, A. (2022). Global ocean wave fields show consistent regional trends between 1980 and 2014 in a multi-product ensemble. *Communications Earth & Environment*, 3(1), 1–16. <https://doi.org/10.1038/s43247-022-00654-9>
- Flooding Analysis Tool. (n.d.). Retrieved August 9, 2023, from <https://sealevel.nasa.gov/flooding-analysis-tool/projected-flooding?>
- Fox-Kemper, B., H.T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S.S. Drijfhout, T.L. Edwards, N.R. Golledge, M. Hemer, R.E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I.S. Nurhati, L. Ruiz, J.-B. Sallée, A.B.A. Slangen, & Y. Yu. (2021). Ocean, Cryosphere and Sea Level Change. In Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, & B. Zhou (Eds.), *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 1211–1362). Cambridge University Press. <https://doi.org/10.1017/9781009157896.011>
- Frederikse, T., Landerer, F., Caron, L., Adhikari, S., Parkes, D., Humphrey, V. W., Dangendorf, S., Hogarth, P., Zanna, L., Cheng, L., & Wu, Y.-H. (2020). The causes of sea level rise since 1900. *Nature*, 584(7821), 393–397. <https://doi.org/10.1038/s41586-020-2591-3>
- Ghanbari, M., Arabi, M., Kao, S.-C., Obeysekera, J., & Sweet, W. (2021). Climate Change and Changes in Compound Coastal-Riverine Flooding Hazard Along the U.S. Coasts. *Earth's Future*, 9(5), e2021EF002055. <https://doi.org/10.1029/2021EF002055>
- Goals Project. (2015). *The Baylands and Climate Change: What We Can Do. Baylands Ecosystem Habitat Goals Science Update 2015 prepared by the San Francisco Bay Area Wetlands Ecosystem Goals project. California State Coastal Conservancy. Oakland, CA.* https://www.sfei.org/sites/default/files/biblio_files/Baylands_Complete_Report.pdf
- Govorcín, M., Bekaert, D.P., Sangha, S. (2022). Towards Continental Vertical Land Maps from InSAR - in anticipation of the OPERA Project Displacement Products. AGU Fall Meeting 2022.

- Govorcin, M., Bekaert, D. P., Hamlington, B. D. (2024). InSAR Vertical Land Motion Rates for California, doi: 10.5281/zenodo.11154177.
- Graham, R. M., & De Boer, A. M. (2013). The Dynamical Subtropical Front. *Journal of Geophysical Research: Oceans*, 118(10), 5676–5685. <https://doi.org/10.1002/jgrc.20408>
- Grant, A. R. R., Wein, A. M., Befus, K. M., Hart, J. F., Frame, M. T., Volentine, R., Barnard, P., & Knudsen, K. L. (2021). Changes in Liquefaction Severity in the San Francisco Bay Area with Sea-Level Rise. 308–317. <https://doi.org/10.1061/9780784483695.030>
- Greene, C. A., Gardner, A. S., Schlegel, N.-J., & Fraser, A. D. (2022). Antarctic calving loss rivals ice-shelf thinning. *Nature*, 609(7929), 948–953. <https://doi.org/10.1038/s41586-022-05037-w>
- Griggs, G. (2005). The impacts of coastal armoring. *Shore and Beach*, 73, 13–22. https://www.researchgate.net/profile/Gary-Griggs/publication/285969581_The_impacts_of_coastal_armoring/links/568fe4b708aee91f69a13733/The-impacts-of-coastal-armoring.pdf
- Griggs, G., & Patsch, K. (2019). The Protection/Hardening of California’s Coast: Times Are Changing. *Journal of Coastal Research*, 35(5), 1051–1061. <https://doi.org/10.2112/JCOASTRES-D-19A-00007.1>
- Griggs, G., Patsch, K., & Savoy, L. (Eds.). (2005). *Living with the Changing California Coast*.
- Griggs, G, Árvai, J, Cayan, D, DeConto, R, Fox, J, Fricker, HA, Kopp, RE, Tebaldi, C, Whiteman, EA (California Ocean Protection Council Science Advisory Team Working Group). (2017). *Rising Seas in California: An Update on Sea-Level Rise Science*. California Ocean Science Trust. http://www.oceansciencetrust.org/wp-content/uploads/2017/04/OST-Sea-Level-Rising-Report-Final_Amended.pdf
- Guza, R. T., & Thornton, E. B. (1982). Swash oscillations on a natural beach. *Journal of Geophysical Research: Oceans*, 87(C1), 483–491. <https://doi.org/10.1029/JC087iC01p00483>
- Haasnoot, M., Kwakkel, J. H., Walker, W. E., & ter Maat, J. (2013). Dynamic adaptive policy pathways: A method for crafting robust decisions for a deeply uncertain world. *Global Environmental Change*, 23(2), 485–498. <https://doi.org/10.1016/j.gloenvcha.2012.12.006>
- Habel, S., Fletcher, C. H., Barbee, M. M., & Fornace, K. L. (2024). Hidden Threat: The Influence of Sea-Level Rise on Coastal Groundwater and the Convergence of Impacts on Municipal Infrastructure. *Annual Review of Marine Science*, 16(1), null. <https://doi.org/10.1146/annurev-marine-020923-120737>
- Hamlington, B. D., Chambers, D. P., Frederikse, T., Dangendorf, S., Fournier, S., Buzzanga, B., & Nerem, R. S. (2022). Observation-based trajectory of future sea level for the coastal United States tracks near high-end model projections. *Communications Earth & Environment*, 3(1), 1–11. <https://doi.org/10.1038/s43247-022-00537-z>
- Hamlington, B. D., Frederikse, T., Thompson, P. R., Willis, J. K., Nerem, R. S., & Fasullo, J. T. (2021). Past, Present, and Future Pacific Sea-Level Change. *Earth’s Future*, 9(4), e2020EF001839. <https://doi.org/10.1029/2020EF001839>
- Hamlington, B. D., Osler, M., Vinogradova, N., & Sweet, W. V. (2021). Coordinated Science Support for Sea-Level Data and Services in the United States. *AGU Advances*, 2(2), e2021AV000418. <https://doi.org/10.1029/2021AV000418>
- Hammond, W. C., Blewitt, G., Kreemer, C., & Nerem, R. S. (2021). GPS Imaging of Global Vertical Land Motion for Studies of Sea Level Rise. *Journal of Geophysical Research: Solid Earth*, 126(7), e2021JB022355. <https://doi.org/10.1029/2021JB022355>

- Hapke, C. J., & Reid, D. (2007). National Assessment of Shoreline Change, Part 4: Historical Coastal Cliff Retreat along the California Coast. U.S. Geological Survey Open-file Report 2007-1133. U.S. Geological Survey <https://pubs.usgs.gov/of/2007/1133/>
- Heady, W. N., Cohen, B. S., Gleason, M. G., Morris, J. N., Newkirk, S. G., Klausmeyer, K. R., Walecka, H., Gagneron, E., Small, M. (2018). Conserving California's Coastal Habitats: A Legacy and a Future with Sea Level Rise. The Nature Conservancy. San Francisco, CA. California State Coastal Conservancy, Oakland, CA. 143 pages. https://conservationgateway.org/ConservationPractices/Marine/crr/library/Documents/TNC_SCC_CoastalAssessment_lo%20snl.pdf
- Hill, K., Hirschfeld, D., Lindquist, C., Cook, F., & Warner, S. (2023). Rising Coastal Groundwater as a Result of Sea-Level Rise Will Influence Contaminated Coastal Sites and Underground Infrastructure. *Earth's Future*, 11(9), e2023EF003825. <https://doi.org/10.1029/2023EF003825>
- Hino, M., Field, C. B., & Mach, K. J. (2017). Managed retreat as a response to natural hazard risk. *Nature Climate Change*, 7(5), 364-370. <https://doi.org/10.1038/nclimate3252>
- Hirschfeld, D., & Hill, K. E. (2022). The landscape of sea level rise adaptation resources: Applying grounded theory in California. *Climate Services*, 28, 100332. <https://doi.org/10.1016/j.cliser.2022.100332>
- Hoover, D. J., Odigie, K. O., Swarzenski, P. W., & Barnard, P. (2017). Sea-level rise and coastal groundwater inundation and shoaling at select sites in California, USA. *Journal of Hydrology: Regional Studies*, 11, 234-249. <https://doi.org/10.1016/j.ejrh.2015.12.055>
- Hu, Y., & Fu, Q. (2007). Observed poleward expansion of the Hadley circulation since 1979. *Atmospheric Chemistry and Physics*, 7(19), 5229-5236. <https://doi.org/10.5194/acp-7-5229-2007>
- Interagency Sea Level Rise Scenario Tool. (n.d.). NASA Sea Level Change Portal. Retrieved August 9, 2023, from <https://sealevel.nasa.gov/task-force-scenario-tool>
- Jakovovic, D., Werner, A. D., de Louw, P. G. B., Post, V. E. A., & Morgan, L. K. (2016). Saltwater upconing zone of influence. *Advances in Water Resources*, 94, 75-86. <https://doi.org/10.1016/j.advwatres.2016.05.003>
- Jones, J. M., Henry, K., Wood, N., Ng, P., & Jamieson, M. (2017). HERA: A dynamic web application for visualizing community exposure to flood hazards based on storm and sea level rise scenarios. *Computers & Geosciences*, 109, 124-133. <https://doi.org/10.1016/j.cageo.2017.08.012>
- Judge, J., Newkirk, S., Leo, K., Heady, W., Hayden, M., Veloz, S., Cheng, T., Battalio, B., Ursell, T., & Small, M. (2017). Case Studies of Natural Shoreline Infrastructure in Coastal California: A Component of Identification of Natural Infrastructure Options for Adapting to Sea Level Rise (California's Fourth Climate Change Assessment). The Nature Conservancy. http://scc.ca.gov/files/2017/11/tnc_Natural-Shoreline-Case-Study_hi.pdf
- Kauffman, N., & Hill, K. (2021). Climate Change, Adaptation Planning and Institutional Integration: A Literature Review and Framework. *Sustainability*, 13(19), 10708. <https://doi.org/10.3390/su131910708>
- Kirwan, M. L., & Megonigal, J. P. (2013). Tidal wetland stability in the face of human impacts and sea level rise. *Nature*, 504(7478), 53-60. <https://doi.org/10.1038/nature12856>
- Kopp, R. E., Garner, G. G., Hermans, T. H. J., Jha, S., Kumar, P., Slangen, A. B. A., Turilli, M., Edwards, T. L., Gregory, J. M., Koubbe, G., Levermann, A., Merzky, A., Nowicki, S., Palmer, M. D., & Smith, C. (2023). The Framework for Assessing Changes To Sea-level (FACTS) v1.0-rc: A platform for characterizing parametric and structural uncertainty in future global, relative, and extreme sea level change. *EGUsphere*, 1-34. <https://doi.org/10.5194/egusphere-2023-14>

- Leonard, L. A. (1997). Controls of sediment transport and deposition in an incised mainland marsh basin, southeastern North Carolina. *Wetlands*, 17(2), 263–274. <https://doi.org/10.1007/BF03161414>
- Lester, C., Manley, C., Dinh, Y., Rozal, S., Cooper, A., Winters, L., Munster, K., Bok, T., & Wrubel, N. (2023). Planning for Sea Level Rise on California’s Coast: Status, Trends, and Recommendations. Ocean and Coastal Policy Center, Marine Science Institute, University of California, Santa Barbara.
- Limber, P. W., Barnard, P. L., Vitousek, S., & Erikson, L. H. (2018). A Model Ensemble for Projecting Multidecadal Coastal Cliff Retreat During the 21st Century. *Journal of Geophysical Research: Earth Surface*, 123(7), 1566–1589. <https://doi.org/10.1029/2017JF004401>
- Loáiciga, H. A., Pingel, T. J., & Garcia, E. S. (2012). Sea Water Intrusion by Sea-Level Rise: Scenarios for the 21st Century. *Groundwater*, 50(1), 37–47. <https://doi.org/10.1111/j.1745-6584.2011.00800.x>
- Longuet-Higgins, M. S., & Stewart, R. w. (1964). Radiation stresses in water waves; a physical discussion, with applications. *Deep Sea Research and Oceanographic Abstracts*, 11(4), 529–562. [https://doi.org/10.1016/0011-7471\(64\)90001-4](https://doi.org/10.1016/0011-7471(64)90001-4)
- Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, & B. Zhou (Eds.). (2021). *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press. <https://www.ipcc.ch/report/ar6/wg1/>
- May, C., Mohan, A., Plane, E., Lopez, D., Mak, M., Luchinsky, L., Hale, A., & Hill, K. (2022). Shallow Groundwater Response to Sea-Level Rise: Alameda, Marin, San Francisco, and San Mateo Counties. <https://doi.org/10.13140/RG.2.2.16973.72164>
- Meucci, A., Young, I. R., Hemer, M., Kirezci, E., & Ranasinghe, R. (2020). Projected 21st century changes in extreme wind-wave events. *Science Advances*, 6(24), eaaz7295. <https://doi.org/10.1126/sciadv.aaz7295>
- Michael, H. A., Russoniello, C. J., & Byron, L. A. (2013). Global assessment of vulnerability to sea level rise in topography-limited and recharge-limited coastal groundwater systems. *Water Resources Research*, 49(4), 2228–2240. <https://doi.org/10.1002/wrcr.20213>
- Mitrovica, J. X., Gomez, N., Morrow, E., Hay, C., Latychev, K., & Tamisiea, M. E. (2011). On the robustness of predictions of sea level fingerprints. *Geophysical Journal International*, 187(2), 729–742. <https://doi.org/10.1111/j.1365-246X.2011.05090.x>
- Moftakhari, H. R., AghaKouchak, A., Sanders, B. F., & Matthew, R. A. (2017). Cumulative hazard: The case of nuisance flooding. *Earth’s Future*, 5(2), 214–223. <https://doi.org/10.1002/2016EF000494>
- Monterey County Water Resources Agency. (2013). Protective Elevations to Control Sea Water Intrusion in the Salinas Valley. Technical Memorandum. https://digitalcommons.csUMB.edu/cgi/viewcontent.cgi?article=1019&context=hornbeck_cgb_5
- Monthly Outlook. (n.d.). Retrieved August 9, 2023, from <https://tidesandcurrents.noaa.gov/high-tide-flooding/monthly-outlook.html>
- Moon, T., Joughin, I., & Smith, B. (2015). Seasonal to multiyear variability of glacier surface velocity, terminus position, and sea ice/ice mélange in northwest Greenland. *Journal of Geophysical Research: Earth Surface*, 120(5), 818–833. <https://doi.org/10.1002/2015JF003494>

- Myers, M. R., Barnard, P. L., Beighley, E., Cayan, D. R., Dugan, J. E., Feng, D., Hubbard, D. M., Iacobellis, S. F., Melack, J. M., & Page, H. M. (2019). A multidisciplinary coastal vulnerability assessment for local government focused on ecosystems, Santa Barbara area, California. *Ocean & Coastal Management*, 182, 104921. <https://doi.org/10.1016/j.ocecoaman.2019.104921>
- Nerem, R. S., Beckley, B. D., Fasullo, J. T., Hamlington, B. D., Masters, D., & Mitchum, G. T. (2018). Climate-change-driven accelerated sea level rise detected in the altimeter era. *Proceedings of the National Academy of Sciences*, 115(9), 2022-2025. <https://doi.org/10.1073/pnas.1717312115>
- Nerem, R. S., Frederikse, T., & Hamlington, B. D. (2022). Extrapolating Empirical Models of Satellite-Observed Global Mean Sea Level to Estimate Future Sea Level Change. *Earth's Future*, 10(4), e2021EF002290. <https://doi.org/10.1029/2021EF002290>
- Newkirk, Sarah, Sam Veloz, Maya Hayden, Walter Heady, Kelly Leo, Jenna Judge, Robert Battalio, Tiffany Cheng, Tara Ursell, & Mary Small. (The Nature Conservancy and Point Blue Conservation Science). (2018). *Toward Natural Infrastructure to Manage Shoreline Change in California* (Publication No. CCCA4-CNRA-2018-011). California's Fourth Climate Change Assessment, California Natural Resources Agency. https://www.energy.ca.gov/sites/default/files/2019-12/Oceans_CCCA4-CNRA-2018-011_ada.pdf
- NOAA | Coastal County Snapshots. (n.d.). Retrieved August 9, 2023, from <https://coast.noaa.gov/snapshots/>
- NOAA. The National Significance of California's Ocean Economy. NOAA Office for Coastal Management. <https://coast.noaa.gov/data/digitalcoast/pdf/california-ocean-economy.pdf>
- Nyman, J. A., Walters, R. J., Delaune, R. D., & Patrick, W. H. (2006). Marsh vertical accretion via vegetative growth. *Estuarine, Coastal and Shelf Science*, 69(3), 370-380. <https://doi.org/10.1016/j.ecss.2006.05.041>
- Ohenhen, L. O., Shirzaei, M., Ojha, C., Sherpa, S. F., & Nicholls, R. J. (2024). Disappearing cities on US coasts. *Nature*, 627(8002), 108-115. <https://doi.org/10.1038/s41586-024-07038-3>
- Our Coast, Our Future. (n.d.). Retrieved August 9, 2023, from <https://ourcoastourfuture.org/>
- Parkinson, B., Patzschke, C. F., Nikolis, D., Raman, S., & Hellgardt, K. (2021). Molten salt bubble columns for low-carbon hydrogen from CH₄ pyrolysis: Mass transfer and carbon formation mechanisms. *Chemical Engineering Journal*, 417, 127407. <https://doi.org/10.1016/j.cej.2020.127407>
- Patsch, K., & Reineman, D. R. (2023, accepted). Sea-level rise impacts on coastal access. *Shore & Beach*.
- Peltier, W. R. (2004). GLOBAL GLACIAL ISOSTASY AND THE SURFACE OF THE ICE-AGE EARTH: The ICE-5G (VM2) Model and GRACE. *Annual Review of Earth and Planetary Sciences*, 32(1), 111-149. <https://doi.org/10.1146/annurev.earth.32.082503.144359>
- Pennell, K. G., Scammell, M. K., McClean, M. D., Ames, J., Weldon, B., Friguglietti, L., Suuberg, E. M., Shen, R., Indeglia, P. A., & Heiger-Bernays, W. J. (2013). Sewer Gas: An Indoor Air Source of PCE to Consider During Vapor Intrusion Investigations. *Groundwater Monitoring & Remediation*, 33(3), 119-126. <https://doi.org/10.1111/gwmr.12021>
- Plane, E., Hill, K., & May, C. (2019). A Rapid Assessment Method to Identify Potential Groundwater Flooding Hotspots as Sea Levels Rise in Coastal Cities. *Water*, 11(11), 2228. <https://doi.org/10.3390/w11112228>
- Program, U. S. G. C. R. (2023). Fifth National Climate Assessment. In *Fifth National Climate Assessment* (pp. 1-470). U.S. Global Change Research Program, Washington, DC. <https://nca2023.globalchange.gov/chapter/focus-on-1/>

- Rahimi, R., Tavakol-Davani, H., Graves, C., Gomez, A., & Fazel Valipour, M. (2020). Compound Inundation Impacts of Coastal Climate Change: Sea-Level Rise, Groundwater Rise, and Coastal Precipitation. *Water*, 12(10), 2776. <https://doi.org/10.3390/w12102776>
- Ranger, N., Reeder, T., & Lowe, J. (2013). Addressing ‘deep’ uncertainty over long-term climate in major infrastructure projects: Four innovations of the Thames Estuary 2100 Project. *EURO Journal on Decision Processes*, 1(3), 233–262. <https://doi.org/10.1007/s40070-013-0014-5>
- Reguero, B. G., Losada, I. J., & Méndez, F. J. (2019). A recent increase in global wave power as a consequence of oceanic warming. *Nature Communications*, 10(1), 205. <https://doi.org/10.1038/s41467-018-08066-0>
- Reineman, D. R., Thomas, L. N., & Caldwell, M. R. (2017). Using local knowledge to project sea level rise impacts on wave resources in California. *Ocean & Coastal Management*, 138, 181–191. <https://doi.org/10.1016/j.ocecoaman.2017.01.020>
- Ribal, A., & Young, I. R. (2019). 33 years of globally calibrated wave height and wind speed data based on altimeter observations. *Scientific Data*, 6(1), 77. <https://doi.org/10.1038/s41597-019-0083-9>
- Roghani, M., Li, Y., Rezaei, N., Robinson, A., Shirazi, E., & Pennell, K. G. (2021). Modeling Fate and Transport of Volatile Organic Compounds (VOCs) Inside Sewer Systems. *Groundwater Monitoring & Remediation*, 41(2), 112–121. <https://doi.org/10.1111/gwmmr.12449>
- S.B. 1, 2022 Biennium, 2021 S. Sess. (Cal 2021). https://leginfo.legislature.ca.gov/faces/billTextClient.xhtml?bill_id=202120220SB1
- S.B. 272, 2016 Biennium, 2015 S. Sess. (Cal 2015). https://leginfo.legislature.ca.gov/faces/billTextClient.xhtml?bill_id=201520160SB272
- Sadrpour, N., & Reineman, D. R. (2023). The impacts of climate change on surfing resources. *Shore & Beach*, 91(1), 32–48. <https://doi.org/10.34237/1009113>
- San Diego Regional Climate Collaborative—School of Leadership and Education Sciences—University of San Diego. (n.d.). Retrieved August 9, 2023, from <https://www.sandiego.edu/soles/centers-and-institutes/nonprofit-institute/signature-programs/climate-collaborative>
- Sea Level Rise and Coastal Flooding Impacts. (n.d.). Retrieved August 9, 2023, from <https://coast.noaa.gov/slr/>
- Sea-Level Rise Collaborative. (2022). State Agency Sea-Level Rise Action Plan for California. https://www.opc.ca.gov/webmaster/_media_library/2022/08/SLR-Action-Plan-2022-508.pdf
- See your local sea level and coastal flood risk. (n.d.). Climate Central. Retrieved August 9, 2023, from <http://riskfinder.climatecentral.org>
- Serafin, K. A., Ruggiero, P., & Stockdon, H. F. (2017). The relative contribution of waves, tides, and nontidal residuals to extreme total water levels on U.S. West Coast sandy beaches. *Geophysical Research Letters*, 44(4), 1839–1847. <https://doi.org/10.1002/2016GL071020>
- SFEI and SPUR. (2019). San Francisco Bay Shoreline Adaptation Atlas: Working with Nature to Plan for Sea Level Rise Using Operational Landscape Units. Publication #915, San Francisco Estuary Institute, Richmond, CA. <https://www.sfei.org/documents/adaptationatlas>
- Shirzaei, M., Freymueller, J., Törnqvist, T. E., Galloway, D. L., Dura, T., & Minderhoud, P. S. J. (2021). Measuring, modelling and projecting coastal land subsidence. *Nature Reviews Earth & Environment*, 2(1), 40–58. <https://doi.org/10.1038/s43017-020-00115-x>

- Slagel, M. J., & Griggs, G. B. (2008). Cumulative Losses of Sand to the California Coast by Dam Impoundment. *Journal of Coastal Research*, 24(3 (243)), 571-584. <https://doi.org/10.2112/06-0640.1>
- Smallegan, S. M., Irish, J. L., & van Dongeren, A. R. (2017). Developed barrier island adaptation strategies to hurricane forcing under rising sea levels. *Climatic Change*, 143(1), 173-184. <https://doi.org/10.1007/s10584-017-1988-y>
- Steps to Resilience Overview | U.S. Climate Resilience Toolkit. (n.d.). Retrieved August 9, 2023, from <https://toolkit.climate.gov/steps-to-resilience/steps-resilience-overview>
- Stevenson, S. L. (2012). Significant changes to ENSO strength and impacts in the twenty-first century: Results from CMIP5. *Geophysical Research Letters*, 39(17). <https://doi.org/10.1029/2012GL052759>
- Stokes, C. R., Abram, N. J., Bentley, M. J., Edwards, T. L., England, M. H., Foppert, A., Jamieson, S. S. R., Jones, R. S., King, M. A., Lenaerts, J. T. M., Medley, B., Miles, B. W. J., Paxman, G. J. G., Ritz, C., van de Flierdt, T., & Whitehouse, P. L. (2022). Response of the East Antarctic Ice Sheet to past and future climate change. *Nature*, 608(7922), 275-286. <https://doi.org/10.1038/s41586-022-04946-0>
- Sweet, W.V., B.D. Hamlington, R.E. Kopp, C.P. Weaver, P.L. Barnard, D. Bekaert, W. Brooks, M. Craghan, G. Dusek, T. Frederikse, G. Garner, A.S. Genz, J.P. Krasting, E. Larour, D. Marcy, J.J. Marra, J. Obeysekera, M. Osler, M. Pendleton, D. Roman, L. Schmied, W. Veatch, K.D. White, & C. Zuzak. (2022). Global and Regional Sea Level Rise Scenarios for the United States: Updated Mean Projections and Extreme Water Level Probabilities Along U.S. Coastlines. NOAA Technical Report NOS 01. National Oceanic and Atmospheric Administration, National Ocean Service. <https://oceanservice.noaa.gov/hazards/sealevelrise/noaa-nostechrpt01-global-regional-SLR-scenarios-US.pdf>
- Swirad, Z. M., & Young, A. P. (2022). Spatial and temporal trends in California coastal cliff retreat. *Geomorphology*, 412, 108318. <https://doi.org/10.1016/j.geomorph.2022.108318>
- Taherkhani, M., Vitousek, S., Barnard, P. L., Frazer, N., Anderson, T. R., & Fletcher, C. H. (2020). Sea-level rise exponentially increases coastal flood frequency. *Scientific Reports*, 10(1), 6466. <https://doi.org/10.1038/s41598-020-62188-4>
- Takekawa, J. Y., Woo, I., Gardiner, R., Casazza, M., Ackerman, J. T., Nur, N., Liu, L., & Spautz, H. (2011). Avian Communities in Tidal Salt Marshes of San Francisco Bay: A Review of Functional Groups by Foraging Guild and Habitat Association. *San Francisco Estuary and Watershed Science*, 9(3). <https://doi.org/10.15447/sfew.2011v9iss3art4>
- Tansel, B., & Zhang, K. (2022). Effects of saltwater intrusion and sea level rise on aging and corrosion rates of iron pipes in water distribution and wastewater collection systems in coastal areas. *Journal of Environmental Management*, 315, 115153. <https://doi.org/10.1016/j.jenvman.2022.115153>
- Thompson, P. R., & Mitchum, G. T. (2014). Coherent sea level variability on the North Atlantic western boundary. *Journal of Geophysical Research: Oceans*, 119(9), 5676-5689. <https://doi.org/10.1002/2014JC009999>
- Thompson, P. R., Widlansky, M. J., Hamlington, B. D., Merrifield, M. A., Marra, J. J., Mitchum, G. T., & Sweet, W. (2021). Rapid increases and extreme months in projections of United States high-tide flooding. *Nature Climate Change*, 11(7), 584-590. <https://doi.org/10.1038/s41558-021-01077-8>
- Thorne, K., MacDonald, G., Guntenspergen, G., Ambrose, R., Buffington, K., Dugger, B., Freeman, C., Janousek, C., Brown, L., Rosencranz, J., Holmquist, J., Smol, J., Hargan, K., & Takekawa, J. (2018). U.S. Pacific coastal wetland resilience and vulnerability to sea level rise. *Science Advances*, 4(2), eaao3270. <https://doi.org/10.1126/sciadv.aao3270>

- Toxic Tides and Environmental Injustice: Social Vulnerability to Sea Level Rise and Flooding of Hazardous Sites in Coastal California | Environmental Science & Technology. (n.d.). Retrieved August 9, 2023, from <https://pubs.acs.org/doi/10.1021/acs.est.2c07481>
- US Department of Commerce, N. O. and A. A. (n.d.). NOAA/NOS Vertical Datums Transformation. Retrieved May 13, 2024, from <https://vdatum.noaa.gov/>
- USGS - HERA. (n.d.). Retrieved August 9, 2023, from <https://www.usgs.gov/apps/hera/>
- van de Wal, R. S. W., Nicholls, R. J., Behar, D., McInnes, K., Stammer, D., Lowe, J. A., Church, J. A., DeConto, R., Fettweis, X., Goelzer, H., Haasnoot, M., Haigh, I. D., Hinkel, J., Horton, B. P., James, T. S., Jenkins, A., LeCozannet, G., Levermann, A., Lipscomb, W. H., ... White, K. (2022). A High-End Estimate of Sea Level Rise for Practitioners. *Earth's Future*, 10(11), e2022EF002751. <https://doi.org/10.1029/2022EF002751>
- Vitousek, S., Barnard, P. L., Fletcher, C. H., Frazer, N., Erikson, L., & Storlazzi, C. D. (2017). Doubling of coastal flooding frequency within decades due to sea level rise. *Scientific Reports*, 7(1), 1399. <https://doi.org/10.1038/s41598-017-01362-7>
- Vitousek, S., Cagigal, L., Montaña, J., Rueda, A., Mendez, F., Coco, G., & Barnard, P. L. (2021). The Application of Ensemble Wave Forcing to Quantify Uncertainty of Shoreline Change Predictions. *Journal of Geophysical Research: Earth Surface*, 126(7), e2019JF005506. <https://doi.org/10.1029/2019JF005506>
- Vitousek, S., Vos, K., Splinter, K. D., Erikson, L., & Barnard, P. L. (2023). A Model Integrating Satellite-Derived Shoreline Observations for Predicting Fine-Scale Shoreline Response to Waves and Sea-Level Rise Across Large Coastal Regions. *Journal of Geophysical Research: Earth Surface*, 128(7), e2022JF006936. <https://doi.org/10.1029/2022JF006936>
- Wengel, C., Lee, S.-S., Stuecker, M. F., Timmermann, A., Chu, J.-E., & Schloesser, F. (2021). Future high-resolution El Niño/Southern Oscillation dynamics. *Nature Climate Change*, 11(9), 758-765. <https://doi.org/10.1038/s41558-021-01132-4>
- Werners, S. E., Wise, R. M., Butler, J. R. A., Totin, E., & Vincent, K. (2021). Adaptation pathways: A review of approaches and a learning framework. *Environmental Science & Policy*, 116, 266-275. <https://doi.org/10.1016/j.envsci.2020.11.003>
- Willis, J., Hamlington, B., & Fournier, S. (2023). Global Mean Sea Level, Trajectory and Extrapolation (Version 101) [dataset]. Zenodo. <https://doi.org/10.5281/zenodo.7702315>
- Yu, X., Yang, J., Graf, T., Koneshloo, M., O'Neal, M. A., & Michael, H. A. (2016). Impact of topography on groundwater salinization due to ocean surge inundation. *Water Resources Research*, 52(8), 5794-5812. <https://doi.org/10.1002/2016WR018814>

APPENDIX 1:

Map of California NOAA Tide Gauge Locations



APPENDIX 2:

Sea Level Scenarios at NOAA Tide Gauge Locations

FOR EACH TIDE GAUGE, amounts of sea level rise (in feet) are provided by decade and for each sea level scenario, from 2020 to 2150 (Tables 1 to 14). The results shown in Chapter 2.0, Table 2.1 are statewide averages. Sea Level Scenarios are also produced at each individual NOAA tide gauge location (see Appendix 1), and these incorporate a local estimate of vertical land motion. The difference between the individual tide gauge numbers and the statewide average (Table 2.1) for any given year or scenario reflects the contribution of vertical land motion in that location. The 2000 baseline is set agnostic of vertical datum, but

the scenarios can be set to specific choices of vertical datum for application to projects. This will involve applying a vertical offset to the scenarios to align the 2000 baseline with the baselines of existing tidal datums of interest (e.g. MSL, MHHW). The 2022 Sea Level Rise Technical Report provides regional offsets for 1992-2000, 2000-2005, and 2005-2020 that can be used for these purposes. These scenarios, now set in the appropriate tidal datum, can then be converted to land-based heights (i.e. transform to a geodetic datum such as NAVD88).

TABLE 1. Sea Level Scenarios for Crescent City.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| 2030 | 0.1 | 0.1 | 0.2 | 0.2 | 0.2 |
| 2040 | 0.1 | 0.2 | 0.2 | 0.3 | 0.4 |
| 2050 | 0.1 | 0.3 | 0.4 | 0.6 | 0.8 |
| 2060 | 0.1 | 0.4 | 0.6 | 1.0 | 1.5 |
| 2070 | 0.2 | 0.4 | 0.8 | 1.6 | 2.3 |
| 2080 | 0.2 | 0.6 | 1.2 | 2.3 | 3.4 |
| 2090 | 0.2 | 0.7 | 1.7 | 3.0 | 4.5 |
| 2100 | 0.2 | 0.8 | 2.3 | 3.9 | 5.6 |
| 2110 | 0.2 | 0.9 | 2.9 | 4.7 | 6.9 |
| 2120 | 0.2 | 1.0 | 3.4 | 5.3 | 7.9 |
| 2130 | 0.2 | 1.2 | 3.8 | 5.8 | 8.7 |
| 2140 | 0.2 | 1.3 | 4.2 | 6.3 | 9.6 |
| 2150 | 0.2 | 1.4 | 4.7 | 6.8 | 10.3 |

TABLE 2. Sea Level Scenarios for N. Spit, Humboldt Bay.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 |
| 2030 | 0.5 | 0.6 | 0.6 | 0.6 | 0.7 |
| 2040 | 0.7 | 0.8 | 0.9 | 1 | 1.1 |
| 2050 | 0.9 | 1 | 1.2 | 1.4 | 1.6 |
| 2060 | 1.1 | 1.3 | 1.5 | 2 | 2.4 |
| 2070 | 1.3 | 1.5 | 1.9 | 2.7 | 3.5 |
| 2080 | 1.4 | 1.8 | 2.5 | 3.6 | 4.7 |
| 2090 | 1.6 | 2.1 | 3.1 | 4.5 | 6 |
| 2100 | 1.8 | 2.4 | 3.9 | 5.5 | 7.3 |
| 2110 | 1.9 | 2.7 | 4.6 | 6.5 | 8.7 |
| 2120 | 2.1 | 3 | 5.3 | 7.3 | 9.9 |
| 2130 | 2.3 | 3.3 | 5.9 | 8 | 10.8 |
| 2140 | 2.4 | 3.5 | 6.5 | 8.6 | 11.9 |
| 2150 | 2.6 | 3.8 | 7.1 | 9.3 | 12.8 |

TABLE 3. Sea Level Scenarios for Arena Cove.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| 2030 | 0.2 | 0.3 | 0.3 | 0.4 | 0.4 |
| 2040 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| 2050 | 0.4 | 0.5 | 0.7 | 0.9 | 1.1 |
| 2060 | 0.5 | 0.7 | 0.9 | 1.4 | 1.8 |
| 2070 | 0.5 | 0.8 | 1.2 | 2.1 | 2.8 |
| 2080 | 0.6 | 1.0 | 1.7 | 2.8 | 3.9 |
| 2090 | 0.7 | 1.2 | 2.2 | 3.6 | 5.1 |
| 2100 | 0.8 | 1.4 | 2.9 | 4.5 | 6.4 |
| 2110 | 0.8 | 1.6 | 3.6 | 5.4 | 7.6 |
| 2120 | 0.9 | 1.7 | 4.1 | 6.1 | 8.7 |
| 2130 | 0.9 | 1.9 | 4.6 | 6.7 | 9.6 |
| 2140 | 1.0 | 2.1 | 5.1 | 7.3 | 10.5 |
| 2150 | 1.0 | 2.3 | 5.6 | 7.8 | 11.4 |

TABLE 4. Sea Level Scenarios for Port Chicago.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 |
| 2030 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 |
| 2040 | 0.4 | 0.5 | 0.6 | 0.7 | 0.8 |
| 2050 | 0.5 | 0.6 | 0.8 | 1.0 | 1.3 |
| 2060 | 0.6 | 0.8 | 1.1 | 1.5 | 2.0 |
| 2070 | 0.7 | 1.0 | 1.4 | 2.2 | 2.9 |
| 2080 | 0.8 | 1.2 | 1.8 | 3.0 | 4.1 |
| 2090 | 0.9 | 1.4 | 2.4 | 3.8 | 5.3 |
| 2100 | 1.0 | 1.6 | 3.1 | 4.8 | 6.5 |
| 2110 | 1.0 | 1.8 | 3.8 | 5.6 | 7.8 |
| 2120 | 1.1 | 2.0 | 4.4 | 6.4 | 9.0 |
| 2130 | 1.2 | 2.2 | 4.9 | 7.0 | 9.9 |
| 2140 | 1.3 | 2.4 | 5.4 | 7.6 | 10.8 |
| 2150 | 1.4 | 2.6 | 6.0 | 8.1 | 11.7 |

TABLE 5. Sea Level Scenarios for Point Reyes.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 |
| 2030 | 0.3 | 0.4 | 0.4 | 0.4 | 0.5 |
| 2040 | 0.4 | 0.5 | 0.6 | 0.7 | 0.8 |
| 2050 | 0.5 | 0.7 | 0.8 | 1.0 | 1.3 |
| 2060 | 0.6 | 0.8 | 1.1 | 1.6 | 2.0 |
| 2070 | 0.7 | 1.0 | 1.4 | 2.2 | 3.0 |
| 2080 | 0.8 | 1.2 | 1.9 | 3.0 | 4.1 |
| 2090 | 0.9 | 1.4 | 2.5 | 3.9 | 5.4 |
| 2100 | 1.0 | 1.6 | 3.1 | 4.8 | 6.6 |
| 2110 | 1.1 | 1.8 | 3.8 | 5.7 | 7.9 |
| 2120 | 1.2 | 2.0 | 4.4 | 6.4 | 9.0 |
| 2130 | 1.2 | 2.2 | 4.9 | 7.0 | 9.9 |
| 2140 | 1.3 | 2.4 | 5.5 | 7.6 | 10.9 |
| 2150 | 1.4 | 2.7 | 6.0 | 8.2 | 11.8 |

TABLE 6. Sea Level Scenarios for San Francisco.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 |
| 2030 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 |
| 2040 | 0.4 | 0.5 | 0.6 | 0.7 | 0.8 |
| 2050 | 0.5 | 0.6 | 0.8 | 1.0 | 1.3 |
| 2060 | 0.6 | 0.8 | 1.1 | 1.5 | 2.0 |
| 2070 | 0.7 | 1.0 | 1.4 | 2.2 | 2.9 |
| 2080 | 0.8 | 1.2 | 1.8 | 3.0 | 4.1 |
| 2090 | 0.9 | 1.4 | 2.4 | 3.8 | 5.3 |
| 2100 | 1.0 | 1.6 | 3.1 | 4.8 | 6.5 |
| 2110 | 1.0 | 1.8 | 3.8 | 5.6 | 7.8 |
| 2120 | 1.1 | 2.0 | 4.4 | 6.4 | 9.0 |
| 2130 | 1.2 | 2.2 | 4.9 | 7.0 | 9.9 |
| 2140 | 1.3 | 2.4 | 5.4 | 7.6 | 10.8 |
| 2150 | 1.3 | 2.6 | 6.0 | 8.1 | 11.7 |

TABLE 7. Sea Level Scenarios for Alameda.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| 2030 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 |
| 2040 | 0.3 | 0.4 | 0.4 | 0.5 | 0.6 |
| 2050 | 0.3 | 0.5 | 0.6 | 0.9 | 1.1 |
| 2060 | 0.4 | 0.6 | 0.9 | 1.4 | 1.8 |
| 2070 | 0.5 | 0.8 | 1.2 | 2.0 | 2.7 |
| 2080 | 0.5 | 0.9 | 1.6 | 2.8 | 3.8 |
| 2090 | 0.6 | 1.1 | 2.1 | 3.5 | 5.0 |
| 2100 | 0.6 | 1.2 | 2.8 | 4.4 | 6.2 |
| 2110 | 0.7 | 1.4 | 3.4 | 5.3 | 7.5 |
| 2120 | 0.7 | 1.6 | 4.0 | 5.9 | 8.5 |
| 2130 | 0.8 | 1.7 | 4.5 | 6.5 | 9.4 |
| 2140 | 0.8 | 1.9 | 4.9 | 7.1 | 10.3 |
| 2150 | 0.8 | 2.1 | 5.4 | 7.6 | 11.2 |

TABLE 8. Sea Level Scenarios for Monterey.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| 2030 | 0.2 | 0.3 | 0.3 | 0.4 | 0.4 |
| 2040 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| 2050 | 0.4 | 0.6 | 0.7 | 0.9 | 1.2 |
| 2060 | 0.5 | 0.7 | 1.0 | 1.4 | 1.9 |
| 2070 | 0.6 | 0.9 | 1.3 | 2.1 | 2.8 |
| 2080 | 0.6 | 1.0 | 1.7 | 2.9 | 3.9 |
| 2090 | 0.7 | 1.2 | 2.3 | 3.7 | 5.2 |
| 2100 | 0.8 | 1.4 | 2.9 | 4.6 | 6.4 |
| 2110 | 0.8 | 1.6 | 3.6 | 5.5 | 7.7 |
| 2120 | 0.9 | 1.8 | 4.2 | 6.2 | 8.8 |
| 2130 | 0.9 | 1.9 | 4.7 | 6.8 | 9.7 |
| 2140 | 1.0 | 2.1 | 5.2 | 7.3 | 10.6 |
| 2150 | 1.1 | 2.3 | 5.7 | 7.9 | 11.5 |

TABLE 9. Sea Level Scenarios for Port San Luis.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| 2030 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 |
| 2040 | 0.3 | 0.4 | 0.4 | 0.5 | 0.6 |
| 2050 | 0.3 | 0.5 | 0.6 | 0.9 | 1.1 |
| 2060 | 0.4 | 0.6 | 0.9 | 1.4 | 1.8 |
| 2070 | 0.5 | 0.7 | 1.2 | 2.0 | 2.7 |
| 2080 | 0.5 | 0.9 | 1.6 | 2.8 | 3.8 |
| 2090 | 0.5 | 1.1 | 2.1 | 3.5 | 5.0 |
| 2100 | 0.6 | 1.2 | 2.8 | 4.5 | 6.3 |
| 2110 | 0.6 | 1.4 | 3.4 | 5.3 | 7.5 |
| 2120 | 0.7 | 1.5 | 4.0 | 6.0 | 8.6 |
| 2130 | 0.7 | 1.7 | 4.4 | 6.6 | 9.5 |
| 2140 | 0.7 | 1.9 | 4.9 | 7.1 | 10.4 |
| 2150 | 0.8 | 2.0 | 5.5 | 7.6 | 11.3 |

TABLE 10. Sea Level Scenarios for Santa Barbara.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| 2030 | 0.2 | 0.3 | 0.3 | 0.3 | 0.4 |
| 2040 | 0.3 | 0.4 | 0.4 | 0.5 | 0.6 |
| 2050 | 0.3 | 0.5 | 0.6 | 0.9 | 1.1 |
| 2060 | 0.4 | 0.6 | 0.9 | 1.4 | 1.8 |
| 2070 | 0.5 | 0.7 | 1.2 | 2.0 | 2.7 |
| 2080 | 0.5 | 0.9 | 1.6 | 2.8 | 3.8 |
| 2090 | 0.5 | 1.1 | 2.1 | 3.5 | 5.0 |
| 2100 | 0.6 | 1.2 | 2.8 | 4.5 | 6.3 |
| 2110 | 0.6 | 1.4 | 3.4 | 5.3 | 7.5 |
| 2120 | 0.7 | 1.5 | 4.0 | 6.0 | 8.6 |
| 2130 | 0.7 | 1.7 | 4.4 | 6.6 | 9.5 |
| 2140 | 0.7 | 1.9 | 4.9 | 7.1 | 10.4 |
| 2150 | 0.8 | 2.0 | 5.5 | 7.6 | 11.3 |

TABLE 11. Sea Level Scenarios for Santa Monica.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| 2030 | 0.3 | 0.3 | 0.4 | 0.4 | 0.4 |
| 2040 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| 2050 | 0.4 | 0.6 | 0.7 | 0.9 | 1.2 |
| 2060 | 0.5 | 0.7 | 1.0 | 1.5 | 1.9 |
| 2070 | 0.6 | 0.9 | 1.3 | 2.1 | 2.8 |
| 2080 | 0.6 | 1.0 | 1.7 | 2.9 | 3.9 |
| 2090 | 0.7 | 1.2 | 2.3 | 3.7 | 5.2 |
| 2100 | 0.8 | 1.4 | 2.9 | 4.6 | 6.4 |
| 2110 | 0.8 | 1.6 | 3.6 | 5.5 | 7.7 |
| 2120 | 0.9 | 1.8 | 4.2 | 6.2 | 8.8 |
| 2130 | 0.9 | 1.9 | 4.7 | 6.8 | 9.7 |
| 2140 | 1.0 | 2.1 | 5.2 | 7.3 | 10.6 |
| 2150 | 1.1 | 2.3 | 5.7 | 7.9 | 11.5 |

TABLE 12. Sea Level Scenarios for Los Angeles.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| 2030 | 0.2 | 0.3 | 0.3 | 0.4 | 0.4 |
| 2040 | 0.3 | 0.4 | 0.5 | 0.6 | 0.7 |
| 2050 | 0.4 | 0.5 | 0.7 | 0.9 | 1.1 |
| 2060 | 0.4 | 0.6 | 0.9 | 1.4 | 1.8 |
| 2070 | 0.5 | 0.8 | 1.2 | 2.1 | 2.7 |
| 2080 | 0.5 | 0.9 | 1.6 | 2.8 | 3.8 |
| 2090 | 0.6 | 1.1 | 2.2 | 3.6 | 5.0 |
| 2100 | 0.6 | 1.3 | 2.8 | 4.5 | 6.3 |
| 2110 | 0.7 | 1.4 | 3.5 | 5.3 | 7.6 |
| 2120 | 0.7 | 1.6 | 4.0 | 6.0 | 8.6 |
| 2130 | 0.8 | 1.8 | 4.5 | 6.6 | 9.5 |
| 2140 | 0.8 | 1.9 | 5.0 | 7.1 | 10.4 |
| 2150 | 0.8 | 2.1 | 5.5 | 7.7 | 11.3 |

TABLE 13. Sea Level Scenarios for La Jolla.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.3 | 0.3 | 0.3 |
| 2030 | 0.3 | 0.4 | 0.4 | 0.4 | 0.5 |
| 2040 | 0.4 | 0.5 | 0.6 | 0.7 | 0.8 |
| 2050 | 0.5 | 0.7 | 0.8 | 1.0 | 1.3 |
| 2060 | 0.6 | 0.8 | 1.1 | 1.6 | 2.0 |
| 2070 | 0.7 | 1.0 | 1.4 | 2.3 | 3.0 |
| 2080 | 0.8 | 1.2 | 1.8 | 3.1 | 4.1 |
| 2090 | 0.9 | 1.4 | 2.4 | 3.9 | 5.3 |
| 2100 | 0.9 | 1.6 | 3.1 | 4.8 | 6.6 |
| 2110 | 1.0 | 1.8 | 3.8 | 5.7 | 7.9 |
| 2120 | 1.1 | 2.0 | 4.4 | 6.4 | 9.0 |
| 2130 | 1.2 | 2.2 | 4.9 | 7.1 | 9.9 |
| 2140 | 1.2 | 2.4 | 5.5 | 7.6 | 10.9 |
| 2150 | 1.3 | 2.6 | 6.0 | 8.2 | 11.8 |

TABLE 14. Sea Level Scenarios for San Diego.

Median values of Sea Level Scenarios, in feet, for each decade from 2020 to 2150, with a baseline of 2000. All median scenario values incorporate the local estimate of vertical land motion.

| YEAR | LOW | INT-LOW | INTERMEDIATE | INT-HIGH | HIGH |
|------|-----|---------|--------------|----------|------|
| 2020 | 0.2 | 0.2 | 0.3 | 0.3 | 0.3 |
| 2030 | 0.3 | 0.4 | 0.4 | 0.5 | 0.5 |
| 2040 | 0.4 | 0.5 | 0.6 | 0.7 | 0.8 |
| 2050 | 0.5 | 0.7 | 0.8 | 1.1 | 1.3 |
| 2060 | 0.6 | 0.9 | 1.1 | 1.6 | 2.0 |
| 2070 | 0.7 | 1.0 | 1.4 | 2.3 | 3.0 |
| 2080 | 0.8 | 1.2 | 1.9 | 3.1 | 4.1 |
| 2090 | 0.9 | 1.4 | 2.5 | 3.9 | 5.4 |
| 2100 | 1.0 | 1.6 | 3.2 | 4.9 | 6.7 |
| 2110 | 1.1 | 1.8 | 3.9 | 5.7 | 8.0 |
| 2120 | 1.2 | 2.1 | 4.5 | 6.5 | 9.1 |
| 2130 | 1.3 | 2.3 | 5.0 | 7.1 | 10.0 |
| 2140 | 1.3 | 2.5 | 5.6 | 7.7 | 11.0 |
| 2150 | 1.4 | 2.7 | 6.1 | 8.3 | 11.9 |

APPENDIX 3:

Localized Assessment of Vertical Land Motion (VLM) and Incorporation into Sea Level Scenarios

ACCURATE FUTURE PROJECTIONS

of VLM require an understanding of and accounting for the underlying processes and the time and space scales on which they vary. This report's VLM projections are partially derived from historical data analysis. Estimates of VLM rates are provided at tide gauge locations and on 1-degree grids using a statistical model based on tide gauge data up to around 2019.⁸⁴ This model divides tide gauge data into three components: 1) a global sea level rise signal,⁸⁵ 2) a long-term linear—but regionally varying—rate, and 3) local effects that vary in time and by region. It is the second component that establishes the linear VLM rates utilized in this report, which are integrated into the relative sea level (RSL) projections for each global mean sea level rise scenario. These rates are assumed to extend linearly from historical data into future projections, though such persistence may not always be accurate (e.g., changes in groundwater extraction) but is often assumed due to data limitations.

In recent decades, GNSS stations across U.S. coastal areas have offered VLM estimates that serve as a benchmark for the rates discussed in this report. However, these GNSS-based estimates have shorter record lengths compared to the tide gauge data used here, and many tide gauge locations lack corresponding GNSS stations. Despite

these limitations, advancements such as the GNSS-imaging technique⁸⁶ have enabled the estimation of VLM at all tide gauge sites, using the coastal GNSS network. Furthermore, the adoption of satellite-based Interferometric Synthetic Aperture Radar (InSAR) technology offers more detailed VLM measurements. When calibrated against land GNSS station data, InSAR provides high-resolution VLM rates across extensive regions of the U.S. coastal plain.⁸⁷ This enhanced resolution allows for a more precise understanding of VLM at very localized levels (such as the street block), aiding in better-informed decisions regarding future RSL projections. An updated InSAR assessment providing localized rates of VLM is shown below in Figure A3.1.⁸⁸ The InSAR analysis shows very localized rates of uplift and subsidence (e.g. Humboldt Bay), and broader areas of subsidence north of the Bay Area and along the southern coast of California.

The integration of data from tide gauges, GNSS, and InSAR is crucial for accurately assessing VLM rates and their future impact at scales critical to coastal communities. In parallel, an understanding of the physical drivers of VLM on a similarly local scale is needed to project measured rates forward in a way that captures the range of possible future scenarios and human activities (e.g. Shirzaei et al., 2021). While this necessary research continues, an alternative path forward

84. Kopp et al., 2014; Sweet et al., 2017; Fox-Kemper et al., 2021; Garner et al., 2021.

85. Dangendorf et al., 2019.

86. Hammond et al., 2021.

87. Examples include the work of Bekaert et al., 2017; Shirzaei et al., 2021; Buzzanga et al., 2020; Bekaert et al., 2019; Ohenhen et al., 2024.

88. Govorcin et al., 2024.

is used in this report that provides a work-around in cases where the VLM contribution from the projection framework does not adequately represent the localized signal. Table 15 provides estimated VLM rates at the 14 gauges from three different sources: the sea level scenarios introduced in Section 2, the GNSS-imaged approach from Hammond et al. (2021), and InSAR estimated rates from Govorcin et al. (2024) leveraging the Observational Products for End-Users from Remote Sensing Analysis (OPERA) project.

To evaluate the best VLM estimate for a given location and to then integrate an updated rate of VLM into the sea level scenarios, the steps below should be taken:

1. Use the InSAR VLM map shown in Figure A3.1 ([download data here](#)) to make a screening level assessment of localized VLM. If the rate is similar to the Scenario VLM rate at the nearest tide gauge, the Table for that tide gauge in Appendix 2 should be used. Note, a difference in rates of 0.5 inches/decade will lead to a 5 inch or 0.4 ft difference in the sea level scenarios in 2100. When it is unknown or unclear which rate to use, the default approach should be to use the sea level scenario at the nearest Tide Gauge as provided in Tables 1-14.
2. If the rates differ significantly (by more than 0.5 inches/decade, for example) and a local assessment or understanding of the underlying drivers of VLM is available, that understanding should be used to inform the selection of the most appropriate rate of VLM for inclusion in the sea level scenarios. In other words, if a known process is leading to a localized VLM signal consistent with the InSAR or GPS estimate in Table 15 and that process is assumed to be persistent, users should continue to the next step.
3. To add a localized rate of VLM from GNSS or InSAR to the sea level scenario at any given time, the starting point is the state-wide sea level scenarios shown in Table 2.1 of the main report. For any given year, the localized VLM should be added using the following computation:

$$\text{Sea Level Scenario}_{\text{local}}(\text{year}) = \text{Sea Level Scenario}_{\text{state}}(\text{year}) - \text{VLM} * (\text{year}-2000)$$

This assumes the sign convention used in Table 15, where positive values indicate uplift and negative values indicate subsidence. Also, the rates in Table 15 should be converted from inches/decade to feet/year. Note, that this substitution will introduce additional uncertainty into the sea level scenarios.

TABLE 15. Vertical land motion rate estimates in inches/decade for tide gauge locations along California coastline. Negative values indicate subsidence, while positive values indicate uplift. Rates for a given location may differ due to record length, time period, methodology and source data. The Scenario VLM rate refers to the rate of vertical land motion that is produced by the projection framework at the foundation of the sea level scenarios. This rate is used to estimate the vertical land motion contributions that are included in the sea level scenarios in Tables 1-14 of Appendix 2.

| TIDE GAUGE | SCENARIO VLM RATE | GNSS VLM RATE | InSAR VLM RATE |
|--------------------------|-------------------|---------------|----------------|
| Crescent City | 0.9 | 0.9 | Not Available |
| Humboldt Bay, North Spit | -1.0 | -0.2 | -1.4 |
| Arena Cove | 0.2 | -0.3 | 0.1 |
| Port Chicago | -0.2 | -0.2 | -0.2 |
| Point Reyes | -0.1 | -0.5 | -1.0 |
| Alameda | 0.3 | -0.4 | -0.1 |
| San Francisco | -0.1 | -0.3 | -0.1 |
| Monterey | 0.1 | -0.1 | 0.3 |
| Port San Luis | 0.3 | 0.1 | -0.3 |
| Santa Barbara | 0.4 | 0.1 | 1.6 |
| Santa Monica | 0.1 | 0.1 | 0.1 |
| Los Angeles | 0.3 | -0.2 | -0.0 |
| La Jolla | -0.1 | -0.5 | -0.6 |
| San Diego | -0.2 | -0.7 | -0.2 |




FIGURE A3.1. Map of vertical land motion rates (mm/year) from Sentinel-1 over the period from 2016-2023 using InSAR analysis for California. Blue indicates subsidence while red indicates areas of uplift. Uncertainty on the rates and variability in the trends over the record are shown in the insets.

APPENDIX 4:

Tools and Resources to Support Visualization of Sea Level Rise and Coastal Hazards

SEVERAL EXISTING geospatial and data visualization tools can be used to support sea level rise and coastal hazard planning efforts. State and local planners, project managers, and members of the public can leverage these tools to visualize the impacts of future plausible Sea Level Scenarios in concert with coastal hazards such as flooding, erosion, and groundwater rise. These tools can help inform risk and vulnerability assessments so that coastal managers have an understanding of how current populations and infrastructure are likely to be affected by sea level rise, and can help inform future strategies that promote coastal resilience. The tools can also be used for communications efforts to help audiences visualize sea level rise and coastal hazard risks.

The following tools comprise a non-exhaustive list of existing resources publicly available at the time of this report's release. It is possible that additional data visualization tools will become available prior to the next Sea Level Rise Guidance update, so this list should be considered as a starting point for identifying the appropriate data sets and visualization tools for sea level rise and coastal hazard planning. In general, the most detailed tool available for a particular area should be used for planning, though in some cases a suite of tools should be evaluated to get a better picture of the possible risks.

- **Coastal Storm Modeling System (CoSMoS)**⁸⁹ is a model that has been developed by the United States Geological Survey (USGS) to make detailed predictions of storm-induced coastal flooding, erosion, cliff failures, and groundwater hazards over large geographic scales. CoSMoS information can either be downloaded as GIS shapefiles through the USGS ScienceBase-Catalog, or can be accessed, viewed, and downloaded through the Our Coast, Our Future flood mapper.
- **Our Coast, Our Future (OCOF)**⁹⁰ provides a user-friendly web-based tool for viewing all CoSMoS model results. OCOF was developed by Point Blue Conservation Science and USGS Pacific Coastal and Marine Science Center to provide a platform for data visualization, synthesis, and download of all outputs produced from CoSMoS.
- **Hazards Exposure Reporting and Analytics (HERA)**⁹¹ application developed by the CoSMoS team displays estimates of residents, businesses, and infrastructure that could be exposed to CoSMoS coastal hazard projections from storms and under each of the sea level rise scenarios. HERA can help communities understand how natural hazards could impact their land, people, infrastructure, and livelihoods. In doing so HERA provides tools and data to help communities as they plan and prepare for natural hazards.

89. <https://www.usgs.gov/centers/pcmssc/science/coastal-storm-modeling-system-cosmos>

90. <https://ourcoastourfuture.org/>

91. <https://www.usgs.gov/apps/hera/>

- **NOAA Sea-Level Rise Viewer⁹²** is a visualization tool for coastal communities showing the potential flooding impacts from sea level rise and high tides. Photo simulations of how future flooding might impact local landmarks are provided, as well as data related to water depth, connectivity, flood frequency, socio-economic vulnerability, wetland loss and migration, and mapping confidence.
- **NASA Flooding Analysis Tool⁹³** allows users to view sea level observations and assess past high-tide flooding frequency, view future changes in high-tide flooding frequency under the Sea Level Scenarios, and view statistics and inflection points that support decision making. This tool was developed by scientists at the University of Hawaii Sea Level Center with funding from the NASA Sea Level Change Team and is based on the methods of Thompson et al. (2021).
- **NOAA's Adapting Stormwater Management for Coastal Floods⁹⁴** tool can be used by communities to determine how current and future flooding can affect their stormwater systems. The website allows practitioners to generate reports that can be used to display local information about observed and projected flooding impacts to inform planning efforts. Beyond providing an interface for analyzing flood data, this tool also includes planning recommendations for how to prepare stormwater management systems for coastal flooding.
- **NOAA's Monthly High Tide Flooding Outlook⁹⁵** shows when and where above-normal high tides and high-tide flooding may be experienced. High-tide flooding likelihoods are updated on a monthly basis, and are derived from a probabilistic model that incorporates tide predictions, sea level rise trends, and seasonal changes in coastal sea level to predict the potential that higher than normal high tide may exceed established National Ocean Service flood thresholds.
- **Coastal County Snapshots⁹⁶** produced by NOAA can be leveraged to produce printable reports describing sea level rise and special flood hazards on a county scale. Users can see data superimposed on a map or through a graphic interface with all data accessible for each snapshot. By delivering complex, county-level data into easy-to-understand charts and graphics, Coastal County Snapshots can support communication about planning decisions or processes.
- **NASA's Interagency Sea Level Rise Scenario Tool⁹⁷** was developed to make the sea level scenarios updated in the 2022 Federal Sea Level Rise Technical Report publicly accessible. The information in the report and this tool is intended to inform coastal communities and others about current and future sea level rise to help contextualize its effects for decision making purposes. The scenarios presented in this tool formed the basis for the California Sea Level Scenarios described in Chapter 2.0 and presented in Appendix 2 of this report. The Interagency Sea Level Rise Scenario Tool was developed by the Sea Level Rise and Coastal Flood Hazard Scenarios and Tools Interagency Task Force, which is the Task Force that authored the federal Technical Report.

92. <https://coast.noaa.gov/slr/>

93. <https://sealevel.nasa.gov/flooding-analysis-tool/projected-flooding?>

94. <https://coast.noaa.gov/stormwater-floods/>

95. <https://tidesandcurrents.noaa.gov/high-tide-flooding/monthly-outlook.html>

96. <https://coast.noaa.gov/snapshots/>

97. <https://sealevel.nasa.gov/task-force-scenario-tool/>

- **Cal-Adapt⁹⁸** makes scientific projections and analyses available as a basis for understanding local climate risks and resilience options. Cal-Adapt is designed to provide the public, researchers, government agencies and industry stakeholders with tools and data for climate adaptation planning that can build resilience and foster community engagement.
 - **Surging Seas Risk Finder⁹⁹** is a multi-part public web tool that provides local sea level rise and flood risk projections, interactive maps, and exposure tabulations from zip codes and up. The Risk Finder aims to provide citizens, communities and policymakers with easily accessible, science-based, local information that can help users understand and respond to the risks of sea level rise and coastal flooding. This tool was collaboratively developed by a sea level rise group led by Climate Central.
 - **Adapting to Rising Tides (ART) Bay Area Flood Explorer¹⁰⁰** provides a regional-scale illustration of coastal flooding due to specific sea level rise and storm surge scenarios. This tool developed by the San Francisco Bay Conservation and Development Commission is intended to improve sea level rise awareness and preparedness by mapping potential future shoreline inundation of areas.
 - **Coastal Resilience¹⁰¹** is a program led by The Nature Conservancy to examine nature's role in reducing coastal flood risk. The program consists of an approach, a web mapping tool, and a network of practitioners around the world supporting hazard mitigation and climate adaptation planning.
- **California King Tides Project¹⁰²** is an effort led by the California Coastal Commission to help visualize future sea level by observing the highest tides of today. The California King Tides Project website features photos of recent King Tides, information on when to expect King Tides, and educator resources.

98. <https://cal-adapt.org/>

99. <https://riskfinder.climatecentral.org/>

100. <https://explorer.adaptingtorisingtides.org/explorer>

101. <https://maps.coastalresilience.org/>

102. <https://www.coastal.ca.gov/kingtides/>

APPENDIX 5:

Technical Detail on Formation of Sea Level Scenarios

SECTION 2.2 describes the process for forming the California sea level scenarios, following closely from the 2022 Federal Sea Level Rise Technical Report. As described, the starting point is the formation of GMSL target values (or ‘gates’) that span the plausible range of future sea level rise. To determine the trajectory or pathway for arriving at these target values, the SSP-based projections from IPCC AR6 are used. For each SSP, an ensemble of thousands of “samples” of the trajectory of sea level rise from 2020 to 2150 was produced in the IPCC AR6. The associated distribution of these samples for each SSP is shown graphically on the left of Figure 2.2. Each individual distribution spans more than one of the five gates that define the sea level scenarios, and thus each SSP-based projection contributes to more than one of the sea level scenarios. Reproducing a similar figure provided in the 2022 Federal Sea Level Rise Technical Report, figure A5.1 below shows the percentage of samples that comes from each IPCC AR6 sea level projection to build the five sea level scenarios used in this report. For ease of visualization, the SSPs are grouped

into Low, Intermediate and High Emissions for medium confidence projections along with their low confidence counterparts.

As an example interpretation of this figure, the Low sea level scenario generally requires a low emissions pathway, while the Intermediate Low sea level scenario arises from a near-equal combination of low, intermediate, and high emissions pathways. The Intermediate Sea Level Scenario includes low emissions trajectories but is mostly related to high emissions scenarios. In fact, the Intermediate, Intermediate-High, and High scenarios are all heavily driven by high emissions scenarios, and differences between these scenarios are associated predominantly with the possible role and contributions of the low-confidence ice-sheet processes. This also indicates the connection between the sea level scenarios and emissions or warming levels. The Low scenario is much more plausible under a low emissions pathway, while Intermediate and higher scenarios are much more likely to be associated with high emissions pathways, as well as with low-confidence ice-sheet processes.




FIGURE A5.1. Percentage of the contributions from different IPCC AR6 sea level projections to each of the five Sea Level scenarios used in this report: Low, Intermediate-Low, Intermediate, Intermediate-High, and High.

An additional important technical detail in the construction of the sea level scenarios is the establishing of target values or gates specifically in 2100. Since all samples from the IPCC AR6 are required to go through a narrow gate in 2100, the range associated with any sea level scenario in 2100 is forced to be very small. Before and after 2100, the range will increase, reflecting the uncertainty associated with both physical processes as well as the regionalization of the sea level scenarios from their global value. This has certain implications for the information provided in this report:

- The five sea level scenarios are presented without likely ranges. The assessment or application of likely ranges should only be done with consideration of the sea level scenario construction and the time period that is being analyzed. Section 2.3.3 describes these considerations.
- The exceedance values in Table 2.2 are only provided for global mean surface air temperature anomalies from the years 2081–2100 relative to the 1850–1900 climatology. Exceedance probabilities can be assessed for other times, but given the additional likely ranges, the separation (or lack thereof) between neighboring sea level scenarios should be considered in the near term (2050 and before), along with the increasing uncertainty in both the trajectory of future sea level rise and warming beyond 2100. Although not shown here, the exceedance probabilities are similar at those other time periods as shown in Table 2.2.

Photo Credits

PG 49 © Joel Avila, Hawkeye Photography.

PG 52 © LV-Olga, courtesy of Shutterstock, Inc.